

**(19) World Intellectual Property
Organization
International Bureau**



(43) International Publication Date
9 September 2005 (09.09.2005)

PCT

(10) International Publication Number
WO 2005/082864 A1

- (51) **International Patent Classification⁷:** C07D 239/22, 401/06, 401/12, 403/06, 403/12, 405/06, 413/06, 417/06, C07F 9/6512, A61K 31/505, 31/506, 31/662, A61P 11/00

(74) **Common Representative:** BAYER HEALTHCARE AG; Law and Patents, Patents and Licensing, 51368 Leverkusen (DE).

(21) **International Application Number:** PCT/EP2005/001486

(22) **International Filing Date:** 15 February 2005 (15.02.2005)

(25) **Filing Language:** English

(26) **Publication Language:** English

(30) **Priority Data:**
04004314.3 26 February 2004 (26.02.2004) EP

(71) **Applicant (for all designated States except US):** BAYER HEALTHCARE AG [DE/DE]; 51368 Leverkusen (DE).

(72) **Inventors; and**

(75) **Inventors/Applicants (for US only):** GIELEN-HAERTWIG, Heike [DE/DE]; Claire-Waldoff-Str. 23, 40789 Monheim (DE). ALBRECHT, Barbara [DE/DE]; Heidestr. 9, 42289 Wülfrath (DE). KELDENICH, Jörg [DE/DE]; Damaschkeweg 49, 42113 Wuppertal (DE). LI, Volkhart [DE/DE]; Im Wiesengrund 40, 42553 Velbert (DE). PERNERSTORFER, Josef [AT/DE]; Gerresheimer Str. 9, 40721 Hilden (DE). SCHLEMMER, Karl-Heinz [DE/DE]; Wildsteig 22 a, 42113 Wuppertal (DE). TELAN, Leila [US/DE]; Rabenweg 42, 42115 Wuppertal (DE).

(81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

 - with international search report
 - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments


For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

- (54) Title:** 1,4-DIARYL-DIHYDROPYRIMIDIN-2-ONES AND THEIR USE AS HUMAN NEUTROPHIL ELASTASE INHIBITORS



- (57) Abstract:** The invention relates to novel heterocyclic derivatives of the general formula (I), processes for their preparation, and their use in medicaments, especially for the treatment of chronic obstructive pulmonary diseases, acute coronary syndrome, acute myocardial infarction and heart failure development.

1,4-DIARYL-DIHYDROPYRIMIDIN-2-ONES AND THEIR USE AS HUMAN NEUTROPHIL ELASTASE INHIBITORS

The present invention relates to novel heterocyclic derivatives, processes for their preparation, and their use in medicaments, especially for the treatment of chronic obstructive pulmonary diseases, acute coronary syndrome, acute myocardial infarction and heart failure development.

5 The fibrous protein elastin, which comprises an appreciable percentage of all protein content in some tissues, such as the arteries, some ligaments, the lungs and the heart, can be hydrolysed or otherwise destroyed by a select group of enzymes classified as elastases. Human leukocyte elastase (HLE, EC 3.4.21.37), also known as human neutrophil elastase (HNE), is a glycosylated, strongly basic serine protease and is found in the azurophilic granules of human polymorphonuclear
10 leukocytes (PMN). HNE is released from activated PMN and has been implicated causally in the pathogenesis of acute and chronic inflammatory diseases. HNE is capable of degrading a wide range of matrix proteins including elastin and collagen, and in addition to these actions on connective tissue HNE has a broad range of inflammatory actions including upregulation of IL-8 gene expression, oedema formation, mucus gland hyperplasia and mucus hypersecretion. It also
15 acts as a mediator of tissue injury by hydrolysing collagen structures, e.g. in the heart after acute myocardial infarction or during the development of heart failure, thus damaging endothelial cells, promoting extravasation of neutrophils adhering to the endothelium and influencing the adhesion process itself.


Pulmonary diseases where HNE is believed to play a role include lung fibrosis, pneumonia, acute
20 respiratory distress syndrome (ARDS), pulmonary emphysema, including smoking-induced emphysema, chronic obstructive pulmonary diseases (COPD) and cystic fibrosis. In cardiovascular diseases, HNE is involved in the enhanced generation of ischaemic tissue injury followed by myocardial dysfunction after acute myocardial infarction and in the remodelling processes occurring during the development of heart failure. HNE has also been causally implicated in
25 rheumatoid arthritis, atherosclerosis, brain trauma, cancer and related conditions in which neutrophil participation is involved.

Thus, inhibitors of HLE activity can be potentially useful in the treatment of a number of inflammatory diseases, especially of chronic obstructive pulmonary diseases [R.A. Stockley, *Neutrophils and protease/antiprotease imbalance*, Am. J. Respir. Crit. Care 160, S49-S52 (1999)].
30 Inhibitors of HLE activity can also be potentially useful in the treatment of acute myocardial syndrome, unstable angina pectoris, acute myocardial infarction and coronary artery bypass grafts (CABG) [C.P. Tiefenbacher et al., *Inhibition of elastase improves myocardial function after repetitive ischaemia and myocardial infarction in the rat heart*, Eur. J. Physiol. 433, S563-S570

(1997); Dinerman et al., *Increased neutrophil elastase release in unstable angina pectoris and acute myocardial infarction*, J. Am. Coll. Cardiol. 15, 1559-1563 (1990)], of the development of heart failure [S.J. Gilbert et al., *Increased expression of promatrix metalloproteinase-9 and neutrophil elastase in canine dilated cardiomyopathy*, Cardio. Res. 34, S377-S383 (1997)] and of
 5 atherosclerosis [Dollery et al., *Neutrophil elastase in human atherosclerotic plaque*, Circulation 107, 2829-2836 (2003)].

The synthesis of 5-ethoxycarbonyl-1-phenyl-6-methyl-4-(3-nitrophenyl)-3,4-dihydropyrimidin-2(1*H*)-one is described in J. Heterocyclic Chem. 38, 1051 (2001). A pharmacological activity of this compound is not mentioned.

10 The present invention relates to compounds of the general formula (I)



wherein

A represents an aryl or heteroaryl ring,

15 R^1 , R^2 and R^3 independently from each other represent hydrogen, halogen, nitro, cyano, C_1 - C_6 -alkyl, hydroxy or C_1 - C_6 -alkoxy, wherein C_1 - C_6 -alkyl and C_1 - C_6 -alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and C_1 - C_4 -alkoxy,

R^4 represents

20 - C_1 - C_6 -alkyl which can be substituted by up to three radicals independently selected from the group consisting of hydroxy, C_1 - C_6 -alkoxycarbonyl and hydroxycarbonyl,

- C₃-C₈-cycloalkylcarbonyl which can be substituted by up to three radicals independently selected from the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- 5 - C₁-C₆-alkylcarbonyl which is substituted by phenyl-C₁-C₆-alkoxy or phenyl-C₁-C₆-alkoxycarbonyl which for their part, in the phenyl moiety, can be substituted by halogen, C₁-C₆-alkyl, hydroxy, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,
- C₆-C₁₀-arylcarbonyl which is substituted by one, two or three radicals independently selected from the group consisting of halogen, cyano, nitro, C₁-C₆-alkyl, trifluoromethyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, amino, C₁-C₆-alkoxycarbonyl, hydroxycarbonyl and phenyl,
- 10 - C₁-C₆-alkoxycarbonyl which is substituted by one or two radicals independently selected from the group consisting of phenyl-C₁-C₆-alkoxy, phenyl-C₁-C₆-alkoxycarbonyl, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonylamino and 5- or 6-membered heterocyclyl, wherein C₁-C₆-alkoxy is further substituted by C₁-C₆-alkoxycarbonyl or hydroxycarbonyl, and 5- or 6-
- 15 membered heterocyclyl is further substituted by hydroxy, oxo, C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,
- heteroarylcarbonyl which is substituted by one or two radicals independently selected from the group consisting of hydroxy, amino, halogen, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl, and which can additionally be substituted by C₁-C₆-alkyl,
- 20 - mono- or di-C₁-C₆-alkylaminocarbonyl wherein the alkyl moiety or at least one alkyl moiety, respectively, is substituted by C₆-C₁₀-aryl which for its part can be further substituted by up to three radicals independently selected from the group consisting of halogen, cyano, trifluoromethyl, C₁-C₆-alkyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- 25 - C₆-C₁₀-arylaminocarbonyl or *N*-(C₁-C₆-alkyl)-*N*-(C₆-C₁₀-aryl)aminocarbonyl wherein aryl is substituted by one, two or three radicals independently selected from the group consisting of halogen, cyano, trifluoromethyl, C₁-C₆-alkyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl, and wherein alkyl, when present, can be substituted by up to three radicals independently selected from the group consisting
- 30 of hydroxy, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- C₃-C₈-cycloalkylaminocarbonyl or *N*-(C₁-C₆-alkyl)-*N*-(C₃-C₈-cycloalkyl)aminocarbonyl wherein cycloalkyl can be substituted by up to three radicals independently selected from


the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl, and wherein alkyl, when present, can be substituted by up to three radicals independently selected from the group consisting of hydroxy, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,

- 5 - heterocyclcarbonyl which is substituted by one, two or three radicals independently selected from the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl, phenyl-C₁-C₆-alkoxycarbonyl, hydroxycarbonyl, 5- or 6-membered heterocycl, 5- or 6-membered heteroaryl and C₆-C₁₀-aryl, wherein C₁-C₆-alkyl is further substituted by hydroxy, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl or hydroxycarbonyl, and wherein
- 10 C₆-C₁₀-aryl can be further substituted by up to three radicals independently selected from the group consisting of halogen, cyano, trifluoromethyl, C₁-C₆-alkyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- 15 - *N*-(heterocycl)aminocarbonyl wherein heterocycl can be further substituted by up to three radicals independently selected from the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl, hydroxycarbonyl and phenyl-C₁-C₆-alkyl,
- 20 - a group of the formula -C(=O)-NR^a-SO₂-R^b wherein R^a represents hydrogen or C₁-C₆-alkyl, and R^b represents C₁-C₆-alkyl which can be substituted by trifluoromethyl, or R^b represents C₆-C₁₀-aryl which can be substituted by C₁-C₆-alkyl, halogen, cyano, nitro or trifluoromethyl,
- 25 or
- 30 - a group of the formula -P(=O)(OR^c)₂ wherein R^c represents hydrogen or C₁-C₆-alkyl,
- R⁵ represents C₁-C₄-alkyl, which can be substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy, C₁-C₆-alkoxy, C₂-C₆-alkenoxy, C₁-C₆-alkylthio, amino, mono- and di-C₁-C₆-alkylamino, arylamino, hydroxycarbonyl, C₁-C₆-alkoxycarbonyl and the radical -O-C₁-C₄-alkyl-O-C₁-C₄-alkyl,
- R⁶ represents hydrogen, C₁-C₆-alkyl, formyl, aminocarbonyl, mono- or di-C₁-C₄-alkylaminocarbonyl, C₃-C₈-cycloalkylcarbonyl, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, *N*-(C₁-C₄-alkylsulfonyl)-aminocarbonyl, *N*-(C₁-C₄-alkylsulfonyl)-*N*-(C₁-C₄-alkyl)-aminocarbonyl, heteroaryl, heterocycl, heteroarylcarbonyl or heterocyclcarbonyl, wherein C₁-C₆-alkyl, mono- and di-C₁-C₄-alkylaminocarbonyl, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, heteroaryl and heterocycl can be substituted with one to three identical or different

radicals selected from the group consisting of aryl, heteroaryl, hydroxy, C₁-C₄-alkoxy, hydroxycarbonyl, C₁-C₆-alkoxycarbonyl, aminocarbonyl, mono- and di-C₁-C₄-alkylamino-carbonyl, amino, mono- and di-C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, tri-(C₁-C₆-alkyl)-silyl, cyano, *N*-(mono- or di-C₁-C₄-alkylamino-C₁-C₄-alkyl)-aminocarbonyl, *N*-(C₁-C₄-alkoxy-C₁-C₄-alkyl)-aminocarbonyl and halogen,

or

R⁶ represents a moiety of the formula



wherein

R^d is selected from the group consisting of hydrogen and C₁-C₆-alkyl, and

n represents an integer of 1 or 2,

or

R⁶ represents a group of the formula -T-U wherein

T represents a C₁-C₆-alkanediyl or C₂-C₆-alkenediyl group

and

U represents

- C₆-C₁₀-aryl or 5- or 6-membered heteroaryl each of which is substituted by one, two or three radicals independently selected from the group consisting of halogen, C₁-C₆-alkyl, 5- or 6-membered heteroaryl and a group of the formula -V-W wherein V represents a bond or a C₁-C₆-alkanediyl or C₂-C₆-alkenediyl group both of which can be further substituted by C₃-C₈-cycloalkyl, and W represents C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,
- a group of the formula -C(=O)-NR^e-SO₂-R^f wherein R^e represents hydrogen or C₁-C₆-alkyl, and R^f represents C₁-C₆-alkyl which can be substituted by trifluoro-

methyl, or R^f represents C_6-C_{10} -aryl which can be substituted by C_1-C_6 -alkyl, halogen, cyano, nitro or trifluoromethyl,

- a group of the formula $-C(=O)-NR^gR^h$ wherein R^g represents hydrogen or C_1-C_6 -alkyl, and R^h represents C_6-C_{10} -aryl which can be substituted by C_1-C_6 -alkoxycarbonyl or hydroxycarbonyl,

- a group of the formula $-C(=O)-NR^i-OR^k$ wherein R^i and R^k independently from each other represent hydrogen or C_1-C_6 -alkyl,

or

- C_6-C_{10} -arylalkoxy which, in the aryl part, can be substituted by halogen, C_1-C_6 -alkyl, C_1-C_6 -alkoxycarbonyl or hydroxycarbonyl,

or

R^6 represents

- C_3-C_8 -cycloalkyl which can be substituted by up to three radicals independently selected from the group consisting of C_1-C_6 -alkyl, hydroxy, oxo, C_1-C_6 -alkoxycarbonyl and hydroxycarbonyl,
- C_2-C_6 -alkenyl which can be substituted by C_1-C_6 -alkoxycarbonyl or hydroxycarbonyl,
- C_1-C_6 -alkylcarbonyl which is substituted by C_1-C_6 -alkoxycarbonylamino,
- C_1-C_6 -alkoxycarbonyl which is substituted by phenyl- C_1-C_6 -alkoxycarbonyl which for its part, in the phenyl moiety, can be further substituted by halogen, C_1-C_6 -alkyl, C_1-C_6 -alkoxycarbonyl or hydroxycarbonyl,

or

- a group of the formula $-SO_2-R^m$ wherein R^m represents C_1-C_6 -alkyl which can be substituted by trifluoromethyl, or R^m represents C_6-C_{10} -aryl which can be substituted by C_1-C_6 -alkyl, halogen, cyano, nitro, trifluoromethyl, C_1-C_6 -alkoxycarbonyl or hydroxycarbonyl,

- R^7 represents halogen, nitro, cyano, C_1-C_6 -alkyl, hydroxy or C_1-C_6 -alkoxy, wherein C_1-C_6 -alkyl and C_1-C_6 -alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and C_1-C_4 -alkoxy,

and

Y^1 , Y^2 , Y^3 , Y^4 and Y^5 independently from each other represent CH or N, wherein the ring contains either 0, 1 or 2 nitrogen atoms.

The compounds according to this invention can also be present in the form of their salts, hydrates
5 and/or solvates.

Physiologically acceptable salts are preferred in the context of the present invention.

Physiologically acceptable salts according to the invention are non-toxic salts which in general are accessible by reaction of the compounds (I) with an inorganic or organic base or acid conventionally used for this purpose. Non-limiting examples of pharmaceutically acceptable salts of
10 compounds (I) include the alkali metal salts, e.g. lithium, potassium and sodium salts, the alkaline earth metal salts such as magnesium and calcium salts, the quaternary ammonium salts such as, for example, triethyl ammonium salts, acetates, benzene sulphonates, benzoates, dicarbonates, disulphates, ditartrates, borates, bromides, carbonates, chlorides, citrates, dihydrochlorides, fumarates, gluconates, glutamates, hexyl resorcinates, hydrobromides, hydrochlorides, hydroxy-
15 naphthoates, iodides, isothionates, lactates, laurates, malates, maleates, mandelates, mesylates, methylbromides, methylnitrates, methylsulphates, nitrates, oleates, oxalates, palmitates, pantothenates, phosphates, diphosphates, polygalacturonates, salicylates, stearates, sulphates, succinates, tartrates, tosylates, valerates, and other salts used for medicinal purposes.

Hydrates of the compounds of the invention or their salts are stoichiometric compositions of the
20 compounds with water, such as for example hemi-, mono-, or dihydrates.

Solvates of the compounds of the invention or their salts are stoichiometric compositions of the compounds with solvents.

The present invention includes both the individual enantiomers or diastereomers and the corresponding racemates or diastereomeric mixtures of the compounds according to the invention
25 and their respective salts. In addition, all possible tautomeric forms of the compounds described above are included according to the present invention. The diastereomeric mixtures can be separated into the individual isomers by chromatographic processes. The racemates can be resolved into the respective enantiomers either by chromatographic processes on chiral phases or by resolution.

30 In the context of the present invention, the substituents, if not stated otherwise, in general have the following meaning:

Alkyl in general represents a straight-chain or branched saturated hydrocarbon radical having 1 to 6, preferably 1 to 4 carbon atoms. Non-limiting examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec.-butyl, tert.-butyl, pentyl, isopentyl, hexyl, isohexyl. The same applies to radicals such as alkoxy, alkylamino, alkylthio, alkylcarbonyl, alkoxycarbonyl, alkoxy-carbonylamino and the like.

Alkanediyl in general represents a straight-chain or branched divalent alkane radical having 1 to 6, preferably 1 to 4 carbon atoms. Non-limiting examples include 1,2-ethylene, 1,3-propylene, propane-1,2-diyl, propane-2,2-diyl, 1,4-butylene, butane-1,3-diyl, butane-2,4-diyl, pentane-2,4-diyl, 2-methyl-pentane-2,4-diyl.

Alkenediyl in general represents a straight-chain or branched divalent alkene radical having 2 to 6, preferably 2 to 4 carbon atoms, and up to three double bonds. Non-limiting examples include ethene-1,2-diyl, ethene-1,1-diyl, propene-1,1-diyl, propene-1,2-diyl, propene-1,3-diyl, propene-3,3-diyl, propene-2,3-diyl, but-2-ene-1,4-diyl, 1,3-butadiene-1,4-diyl, pent-2-ene-1,4-diyl, hex-2-ene-1,4-diyl.

Alkoxy illustratively and preferably represents methoxy, ethoxy, n-propoxy, isopropoxy, tert.-butoxy, n-pentoxy and n-hexoxy.

Arylalkoxy and phenylalkoxy in general represent a straight-chain or branched alkoxy radical which is substituted with an aryl or a phenyl group, respectively. Non-limiting examples include benzyloxy, naphthylmethoxy, 1-phenylethoxy, 2-phenylethoxy, 2-naphthylethoxy, 3-phenylpropoxy, 4-phenyl-butoxy. The same applies to the radical phenylalkoxycarbonyl.

Alkenoxy illustratively and preferably represents allyloxy, but-2-en-1-oxy, pent-3-en-1-oxy and hex-2-en-1-oxy.

Alkylcarbonyl in general represents a straight-chain or branched hydrocarbon radical having 1 to 6, preferably 1 to 4 carbon atoms which has a carbonyl function at the position of attachment. Non-limiting examples include formyl, acetyl, n-propionyl, n-butyryl, isobutyryl, pivaloyl, n-hexanoyl.

Alkylcarbonylamino in general represents a straight-chain or branched hydrocarbon radical having 1 to 6, preferably 1 to 4 carbon atoms which has a carbonylamino (-CO-NH-) function at the position of attachment and which is bonded to the carbonyl group. Non-limiting examples include formylamino, acetylamino, n-propionylamino, n-butyrylamino, isobutyrylamino, pivaloylamino, n-hexanoylamino.

Alkoxy-carbonyl illustratively and preferably represents methoxy-carbonyl, ethoxy-carbonyl, n-propoxy-carbonyl, isopropoxy-carbonyl, tert.-butoxy-carbonyl, n-pentoxy-carbonyl and n-hexoxy-carbonyl.

5 Alkenoxy-carbonyl illustratively and preferably represents allyloxy-carbonyl, but-2-en-1-oxy-carbonyl, pent-3-en-1-oxy-carbonyl and hex-2-en-1-oxy-carbonyl.

Alkylamino represents an alkylamino radical having one or two (independently selected) alkyl residues, illustratively and preferably representing methylamino, ethylamino, n-propylamino, isopropylamino, tert.-butylamino, n-pentylamino, n-hexylamino, *N,N*-dimethylamino, *N,N*-diethylamino, *N*-ethyl-*N*-methylamino, *N*-methyl-*N*-n-propylamino, *N*-isopropyl-*N*-n-propylamino, *N*-tert.-butyl-*N*-methylamino, *N*-ethyl-*N*-n-pentylamino and *N*-n-hexyl-*N*-methylamino.

10

Alkylaminocarbonyl represents an alkylaminocarbonyl radical having one or two (independently selected) alkyl residues, illustratively and preferably representing methylaminocarbonyl, ethylaminocarbonyl, n-propylaminocarbonyl, isopropylaminocarbonyl, tert.-butylaminocarbonyl, n-pentylaminocarbonyl, n-hexylaminocarbonyl, *N,N*-dimethylaminocarbonyl, *N,N*-diethylaminocarbonyl, *N*-ethyl-*N*-methylaminocarbonyl, *N*-methyl-*N*-n-propylaminocarbonyl, *N*-isopropyl-*N*-n-propylaminocarbonyl, *N*-tert.-butyl-*N*-methylaminocarbonyl, *N*-ethyl-*N*-n-pentylaminocarbonyl and *N*-n-hexyl-*N*-methylaminocarbonyl.

15

Alkylsulfonyl in general represents a straight-chain or branched hydrocarbon radical having 1 to 6, preferably 1 to 4 carbon atoms which has a sulfonyl function at the position of attachment. Non-limiting examples include methylsulfonyl, ethylsulfonyl, n-propylsulfonyl, isopropylsulfonyl, n-butylsulfonyl, tert.-butylsulfonyl.

20

Cycloalkyl in general represents a cyclic saturated hydrocarbon radical having 3 to 8, preferably 3 to 6 carbon atoms. Non-limiting examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. The same applies to radicals such as cycloalkylcarbonyl.

25 Aryl in general represents a mono- to tricyclic aromatic carbocyclic radical having 6 to 14, preferably 6 to 10 carbon atoms, illustratively and preferably representing phenyl, naphthyl and phenanthrenyl. The same applies to radicals such as arylcarbonyl, arylalkoxy and arylaminocarbonyl.

Arylcarbonyl illustratively and preferably represents benzoyl and naphthoyl.

Arylaminocarbonyl illustratively and preferably represents phenylaminocarbonyl and naphthylaminocarbonyl.

30

Heteroaryl per se and in heteroarylcarbonyl in general represents an aromatic mono- or bicyclic radical having 5 to 10 and preferably 5 or 6 ring atoms, and up to 5 and preferably up to 4 heteroatoms selected from the group consisting of S, O and N, illustratively and preferably representing thienyl, furyl, pyrrolyl, thiazolyl, oxazolyl, isothiazolyl, isoxazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, pyridyl, pyrimidyl, pyridazinyl, indolyl, indazolyl, benzofuranyl, benzothienyl, benzothiazolyl, quinoliny, isoquinoliny.

Heteroarylcarbonyl illustratively and preferably represents thienylcarbonyl, furylcarbonyl, pyrrolylcarbonyl, thiazolylcarbonyl, oxazolylcarbonyl, isothiazolylcarbonyl, isoxazolylcarbonyl, imidazolylcarbonyl, pyridylcarbonyl, pyrimidylcarbonyl, pyridazinylcarbonyl, indolylcarbonyl, indazolylcarbonyl, benzofuranylcarbonyl, benzothienylcarbonyl, quinoliny carbonyl, isoquinoliny carbonyl.

Heterocyclyl per se and in heterocyclylcarbonyl in general represents a mono- or polycyclic, preferably mono- or bicyclic, non-aromatic heterocyclic radical having 4 to 10 and preferably 5 to 8 ring atoms, and up to 3 and preferably up to 2 heteroatoms and/or hetero-groups selected from the group consisting of N, O, S, SO and SO₂. The heterocyclyl radicals can be saturated or partially unsaturated. Preference is given to 5- to 8-membered monocyclic saturated heterocyclyl radicals having up to two heteroatoms selected from the group consisting of O, N and S, such as illustratively and preferably tetrahydrofuran-2-yl, 1,3-dioxolan-4-yl, pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, pyrroliny, piperidiny, piperazinyl, morpholiny, thiomorpholiny, perhydroazepiny.

Heterocyclylcarbonyl illustratively and preferably represents tetrahydrofuran-2-carbonyl, 1,3-dioxolan-4-carbonyl, pyrrolidin-1-carbonyl, pyrrolidin-2-carbonyl, pyrrolidin-3-carbonyl, pyrrolin-carbonyl, piperidincarbonyl, morpholincarbonyl, perhydroazepincarbonyl.

Halogen represents fluorine, chlorine, bromine and iodine.

When stated, that Y¹, Y², Y³, Y⁴ and Y⁵ represent CH or N, CH shall also stand for a ring carbon atom, which is substituted with a substituent R³ or R⁷.

A * symbol next to a bond denotes the point of attachment in the molecule.

In another preferred embodiment, the present invention relates to compounds of general formula (I), wherein

A represents an aryl or heteroaryl ring,

R¹, R² and R³ independently from each other represent hydrogen, halogen, nitro, cyano, C₁-C₆-alkyl, hydroxy or C₁-C₆-alkoxy, wherein C₁-C₆-alkyl and C₁-C₆-alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and C₁-C₄-alkoxy,

5 R⁴ represents

- C₁-C₆-alkyl which can be substituted by up to three radicals independently selected from the group consisting of hydroxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- C₃-C₈-cycloalkylcarbonyl which can be substituted by up to three radicals independently selected from the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- 10 - C₆-C₁₀-arylcarbonyl which is substituted by one, two or three radicals independently selected from the group consisting of halogen, cyano, C₁-C₆-alkyl, trifluoromethyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- C₁-C₆-alkoxycarbonyl which is substituted by one or two radicals independently selected from the group consisting of phenyl-C₁-C₆-alkoxy, phenyl-C₁-C₆-alkoxycarbonyl, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonylamino and 5- or 6-membered heterocyclyl, wherein C₁-C₆-alkoxy is further substituted by C₁-C₆-alkoxycarbonyl or hydroxycarbonyl, and 5- or 6-membered heterocyclyl is further substituted by hydroxy, oxo, C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,
- 15 - heteroarylcarbonyl which is substituted by one or two radicals independently selected from the group consisting of hydroxy, amino, halogen, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl, and which can additionally be substituted by C₁-C₆-alkyl,
- mono- or di-C₁-C₆-alkylaminocarbonyl wherein the alkyl moiety or at least one alkyl moiety, respectively, is substituted by C₆-C₁₀-aryl which for its part can be further substituted by up to three radicals independently selected from the group consisting of halogen, cyano, trifluoromethyl, C₁-C₆-alkyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- 25 - heterocyclylcarbonyl which is substituted by one, two or three radicals independently selected from the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl, phenyl-C₁-C₆-alkoxycarbonyl, hydroxycarbonyl, 5- or 6-membered heterocyclyl, 5- or 6-membered heteroaryl and C₆-C₁₀-aryl, wherein C₁-C₆-alkyl is further sub-
- 30

stituted by hydroxy, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl or hydroxycarbonyl, and wherein C₆-C₁₀-aryl can be further substituted by up to three radicals independently selected from the group consisting of halogen, cyano, trifluoromethyl, C₁-C₆-alkyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,

5 or


- a group of the formula -C(=O)-NH-SO₂-R^b wherein R^b represents C₁-C₆-alkyl which can be substituted by trifluoromethyl, or R^b represents C₆-C₁₀-aryl which can be substituted by C₁-C₆-alkyl, halogen, cyano, nitro or trifluoromethyl,

10 R⁵ represents C₁-C₄-alkyl, which can be substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy, C₁-C₆-alkoxy, C₂-C₆-alkenoxy, C₁-C₆-alkylthio, amino, mono- and di-C₁-C₆-alkylamino, arylamino, hydroxycarbonyl, C₁-C₆-alkoxycarbonyl and the radical -O-C₁-C₄-alkyl-O-C₁-C₄-alkyl,

15 R⁶ represents hydrogen, C₁-C₆-alkyl, aminocarbonyl, mono- or di-C₁-C₄-alkylaminocarbonyl, C₃-C₈-cycloalkylcarbonyl, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, *N*-(C₁-C₄-alkylsulfonyl)-aminocarbonyl, *N*-(C₁-C₄-alkylsulfonyl)-*N*-(C₁-C₄-alkyl)-aminocarbonyl, heteroarylcarbonyl or heterocyclylcarbonyl, wherein C₁-C₆-alkyl, mono- and di-C₁-C₄-alkylaminocarbonyl, C₁-C₆-alkylcarbonyl and C₁-C₆-alkoxycarbonyl can be substituted with one to three identical or different radicals selected from the group consisting of aryl, heteroaryl, hydroxy, C₁-C₄-alkoxy, hydroxycarbonyl, C₁-C₆-alkoxycarbonyl, aminocarbonyl, 20 mono- and di-C₁-C₄-alkylaminocarbonyl, amino, mono- and di-C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, *N*-(mono- or di-C₁-C₄-alkylamino-C₁-C₄-alkyl)-aminocarbonyl, *N*-(C₁-C₄-alkoxy-C₁-C₄-alkyl)-aminocarbonyl and halogen,

or

R⁶ represents a moiety of the formula



wherein

R^d is selected from the group consisting of hydrogen and C₁-C₆-alkyl, and

n represents an integer of 1 or 2,

or

R⁶ represents a group of the formula -T-U wherein

T represents a C₁-C₄-alkanediyl or C₂-C₄-alkenediyl group

5 and

U represents

10 • C₆-C₁₀-aryl or 5- or 6-membered heteroaryl each of which is substituted by one, two or three radicals independently selected from the group consisting of halogen, C₁-C₆-alkyl, 5- or 6-membered heteroaryl and a group of the formula -V-W wherein V represents a bond, a C₂-C₆-alkenediyl group or a C₁-C₆-alkanediyl group the latter of which can be further substituted by C₃-C₈-cycloalkyl, and W represents C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,

15 • a group of the formula -C(=O)-NH-SO₂-R^f wherein R^f represents C₁-C₆-alkyl which can be substituted by trifluoromethyl, or R^f represents C₆-C₁₀-aryl which can be substituted by C₁-C₆-alkyl, halogen, cyano, nitro or trifluoromethyl,

or

• a group of the formula -C(=O)-NHR^h wherein R^h represents C₆-C₁₀-aryl which can be substituted by C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,

or

20 R⁶ represents

– C₃-C₈-cycloalkyl which can be substituted by up to three radicals independently selected from the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,

or

25 – C₂-C₆-alkenyl which can be substituted by C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,

R⁷ represents halogen, nitro, cyano, C₁-C₆-alkyl, hydroxy or C₁-C₆-alkoxy, wherein C₁-C₆-alkyl and C₁-C₆-alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and C₁-C₄-alkoxy,

and

- 5 Y¹, Y², Y³, Y⁴ and Y⁵ independently from each other represent CH or N, wherein the ring contains either 0, 1 or 2 nitrogen atoms.

In another particularly preferred embodiment, the present invention relates to compounds of general formula (I), wherein

A represents a phenyl, naphthyl or pyridyl ring,

- 10 R¹, R² and R³ independently from each other represent hydrogen, fluoro, chloro, bromo, nitro, cyano, methyl, ethyl, trifluoromethyl or trifluoromethoxy,

R⁴ represents

- C₁-C₄-alkyl which can be substituted by up to two radicals independently selected from the group consisting of hydroxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- 15 – C₃-C₆-cycloalkylcarbonyl which can be substituted by up to two radicals independently selected from the group consisting of C₁-C₄-alkyl, hydroxy, oxo, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- benzoyl which is substituted by one, two or three radicals independently selected from the group consisting of fluoro, chloro, bromo, cyano, C₁-C₄-alkyl, trifluoromethyl, hydroxy, 20 C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- C₁-C₄-alkoxycarbonyl which is substituted by one or two radicals independently selected from the group consisting of benzyloxy, benzyloxycarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-carbonylamino, pyrrolidinyl, piperidinyl and morpholinyl, wherein C₁-C₄-alkoxy is further substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl, and wherein pyrrolidinyl, 25 piperidinyl and morpholinyl is further substituted by hydroxy, oxo, C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,
- furylcarbonyl, thienylcarbonyl, oxazolylcarbonyl, thiazolylcarbonyl, pyridylcarbonyl or pyrimidinylcarbonyl each of which is substituted by one or two radicals independently selected from the group consisting of hydroxy, amino, fluoro, chloro, bromo, C₁-C₄-alkoxy,

C₁-C₄-alkoxycarbonyl and hydroxycarbonyl, and each of which can additionally be substituted by C₁-C₄-alkyl,

5 — mono- or di-C₁-C₄-alkylaminocarbonyl wherein the alkyl moiety or at least one alkyl moiety, respectively, is substituted by phenyl which for its part can be further substituted by up to three radicals independently selected from the group consisting of fluoro, chloro, bromo, cyano, trifluoromethyl, C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,

10 — tetrahydrofurylcarbonyl, tetrahydropyranylcabonyl, piperidinylcarbonyl, piperazinylcarbonyl or morpholinylcarbonyl each of which is substituted by one or two radicals independently selected from the group consisting of C₁-C₄-alkyl, hydroxy, oxo, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl, benzyloxycarbonyl, hydroxycarbonyl, piperidinyl, morpholinyl, pyridyl and phenyl, wherein C₁-C₄-alkyl is further substituted by hydroxy, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl or hydroxycarbonyl, and wherein phenyl can be further substituted by up to three radicals independently selected from the group consisting of fluoro, chloro, bromo, cyano, trifluoromethyl, C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,

or

20 — a group of the formula -C(=O)-NH-SO₂-R^b wherein R^b represents C₁-C₄-alkyl which can be substituted by trifluoromethyl, or R^b represents phenyl which can be substituted by C₁-C₄-alkyl, fluoro, chloro, bromo, cyano, nitro or trifluoromethyl,


R⁵ represents methyl or ethyl,

25 R⁶ represents hydrogen, C₁-C₆-alkyl, mono- or di-C₁-C₄-alkylaminocarbonyl, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl or heterocyclylcarbonyl, wherein C₁-C₆-alkyl and C₁-C₆-alkoxycarbonyl can be substituted with one to three identical or different radicals selected from the group consisting of hydroxy, C₁-C₄-alkoxy, hydroxycarbonyl, C₁-C₆-alkoxycarbonyl, aminocarbonyl, mono- and di-C₁-C₄-alkylaminocarbonyl, amino, mono- and di-C₁-C₄-alkylamino,

or

R⁶ represents a moiety of the formula

16



wherein

R^d is selected from the group consisting of hydrogen and C_1 - C_4 -alkyl, and

n represents an integer of 1 or 2,

5 or

R^6 represents a group of the formula $-\text{T}-\text{U}$ wherein

T represents a C_1 - C_4 -alkanediyl group

and

U represents

10 • phenyl, furyl, thienyl, oxazolyl, thiazolyl or pyridyl each of which is substituted by one or two radicals independently selected from the group consisting of fluoro, chloro, bromo, C_1 - C_4 -alkyl, thienyl, pyridyl and a group of the formula $-\text{V}-\text{W}$ wherein V represents a bond or a C_1 - C_4 -alkanediyl or C_2 - C_4 -alkenediyl group, and W represents C_1 - C_4 -alkoxycarbonyl or hydroxycarbonyl,

15 • a group of the formula $-\text{C}(=\text{O})-\text{NH}-\text{SO}_2-\text{R}^f$ wherein R^f represents C_1 - C_4 -alkyl which can be substituted by trifluoromethyl, or R^f represents phenyl which can be substituted by C_1 - C_4 -alkyl, fluoro, chloro, bromo, cyano, nitro or trifluoromethyl,

or

20 • a group of the formula $-\text{C}(=\text{O})-\text{NHR}^h$ wherein R^h represents phenyl which can be substituted by C_1 - C_4 -alkoxycarbonyl or hydroxycarbonyl,

or

R^6 represents

- C₃-C₆-cycloalkyl which can be substituted by up to two radicals independently selected from the group consisting of C₁-C₄-alkyl, hydroxy, oxo, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,

or

- 5 – C₂-C₄-alkenyl which is substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,

R⁷ represents halogen, nitro, cyano, trifluoromethyl, trifluoromethoxy, methyl or ethyl,

and

Y¹, Y², Y³, Y⁴ and Y⁵ each represent CH.

10 In another very particularly preferred embodiment, the present invention relates to compounds of general formula (I), wherein

A represents a phenyl or a pyridyl ring,

R¹ and R³ each represent hydrogen,

R² represents fluoro, chloro, bromo, nitro or cyano,

R⁴ represents

- 15 – C₁-C₄-alkyl which can be substituted by up to two radicals independently selected from the group consisting of hydroxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- C₃-C₆-cycloalkylcarbonyl which can be substituted by up to two radicals independently selected from the group consisting of C₁-C₄-alkyl, hydroxy, oxo, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- 20 – benzoyl which is substituted by one, two or three radicals independently selected from the group consisting of fluoro, chloro, bromo, cyano, C₁-C₄-alkyl, trifluoromethyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- C₁-C₄-alkoxycarbonyl which is substituted by one or two radicals independently selected from the group consisting of benzyloxy, benzyloxycarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-
- 25 carbonylamino, pyrrolidinyl, piperidinyl and morpholinyl, wherein C₁-C₄-alkoxy is further substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl, and wherein pyrrolidinyl,

piperidinyl and morpholinyl is further substituted by hydroxy, oxo, C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,

5 – furylcarbonyl, oxazolylcarbonyl, thiazolylcarbonyl or pyridylcarbonyl each of which is substituted by one or two radicals independently selected from the group consisting of hydroxy, amino, fluoro, chloro, bromo, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl, and each of which can additionally be substituted by C₁-C₄-alkyl,

10 – mono- or di-C₁-C₄-alkylaminocarbonyl wherein the alkyl moiety or at least one alkyl moiety, respectively, is substituted by phenyl which for its part can be further substituted by up to three radicals independently selected from the group consisting of fluoro, chloro, bromo, cyano, trifluoromethyl, C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,

15 – piperidinylcarbonyl, piperazinylcarbonyl or morpholinylcarbonyl each of which is substituted by one or two radicals independently selected from the group consisting of C₁-C₄-alkyl, hydroxy, oxo, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl, benzyloxycarbonyl, hydroxycarbonyl, piperidinyl, morpholinyl, pyridyl and phenyl, wherein C₁-C₄-alkyl is further substituted by hydroxy, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl or hydroxycarbonyl, and wherein phenyl can be further substituted by up to three radicals independently selected from the group consisting of fluoro, chloro, bromo, cyano, trifluoromethyl, C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,

20 or


– a group of the formula -C(=O)-NH-SO₂-R^b wherein R^b represents C₁-C₄-alkyl which can be substituted by trifluoromethyl, or R^b represents phenyl which can be substituted by C₁-C₄-alkyl, fluoro, chloro, bromo, cyano, nitro or trifluoromethyl,

R⁵ represents methyl,

25 R⁶ represents hydrogen, C₁-C₄-alkyl, mono- or di-C₁-C₄-alkylaminocarbonyl, C₁-C₄-alkylcarbonyl or C₁-C₄-alkoxycarbonyl, wherein C₁-C₄-alkyl and C₁-C₄-alkoxycarbonyl can be substituted with a radical selected from the group consisting of hydroxy, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl, hydroxycarbonyl, aminocarbonyl, mono- and di-C₁-C₄-alkylamino-carbonyl, amino, mono- and di-C₁-C₄-alkylamino,

30 or

R^6 represents a moiety of the formula



wherein

R^d is selected from the group consisting of hydrogen and methyl,

5 or

R^6 represents a group of the formula -T-U wherein

T represents a -CH₂- group

and

U represents

10 • phenyl, furyl or oxazolyl each of which is substituted by one or two radicals independently selected from the group consisting of fluoro, chloro, bromo, C₁-C₄-alkyl and a group of the formula -V-W wherein V represents a bond, a -CH₂- group or a -CH=CH- group, and W represents C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,

15 • a group of the formula -C(=O)-NH-SO₂-R^f wherein R^f represents C₁-C₄-alkyl which can be substituted by trifluoromethyl, or R^f represents phenyl which can be substituted by C₁-C₄-alkyl, fluoro, chloro, bromo, cyano, nitro or trifluoromethyl,

or

20 • a group of the formula -C(=O)-NHR^h wherein R^h represents phenyl which can be substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,

or

R^6 represents

- C₃-C₆-cycloalkyl which can be substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,

or

- a -CH=CH- group which is substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,

R⁷ represents trifluoromethyl or nitro,

and

- 5 Y¹, Y², Y³, Y⁴ and Y⁵ each represent CH.

In another likewise preferred embodiment, the present invention relates to compounds according to general formula (I), wherein A is phenyl or pyridyl.

In another likewise preferred embodiment, the present invention relates to compounds according to general formula (I), wherein R¹ is hydrogen.


- 10 In another likewise preferred embodiment, the present invention relates to compounds according to general formula (I), wherein R² is cyano, especially wherein A is phenyl or pyridyl and R² is cyano located in para-position relative to the central dihydropyrimidinone ring.

In another likewise preferred embodiment, the present invention relates to compounds according to general formula (I), wherein R³ is hydrogen.

- 15 In another likewise preferred embodiment, the present invention relates to compounds according to general formula (I), wherein R⁵ is methyl.

In another likewise preferred embodiment, the present invention relates to compounds according to general formula (I), wherein R⁷ is trifluoromethyl or nitro, especially wherein R⁷ is trifluoromethyl located in meta-position relative to the central dihydropyrimidinone ring.

- 20 In another likewise particularly preferred embodiment, the present invention relates to compounds of general formula (IA)




wherein

Z represents CH or N, and


R¹, R³, R⁴ and R⁶ have the meaning indicated above.

- 5 The compounds of the present invention, wherein R⁶ is hydrogen, can enolize into the corresponding hydroxyamidines:




In another embodiment, the present invention relates to a process for synthesizing the compounds of general formula (I) or (IA), respectively.

- 10 The compounds of general formula (I) or (IA), respectively, can be synthesized by condensing compounds of general formula (II)




wherein A, R¹ and R² have the meaning indicated above,

with compounds of general formula (III)




5 wherein R⁴ and R⁵ have the meaning indicated above,

and compounds of general formula (IV)



wherein R³, R⁷, and Y¹ to Y⁵ have the meaning indicated above,

10 in the presence of an acid or acid anhydride either in a three-component / one-step reaction or sequentially to give compounds of general formula (IB)



wherein A, R¹ to R⁵, R⁷, and Y¹ to Y⁵ have the meaning indicated above,

optionally followed, in case R⁶ does not represent hydrogen, by reaction of the compounds of general formula (IB) with compounds of general formula (V)



wherein

R^{6*} has the meaning of R⁶ as indicated above, but does not represent hydrogen, and

X represents a leaving group, such as halogen, tosylate, mesylate or sulfate,

in the presence of a base.

- 10 Suitable solvents for the process (II) + (III) + (IV) → (IB) are generally customary organic solvents which do not change under the reaction conditions. These include ethers such as diethyl ether, diisopropyl ether, methyl t-butyl ether, 1,2-dimethoxyethane, dioxan or tetrahydrofuran, ethyl acetate, acetone, acetonitrile, dimethylsulfoxide, dimethylformamide, or alcohols such as methanol, ethanol, n-propanol, isopropanol, n-butanol or t-butanol, or hydrocarbons such as
- 15 pentane, hexane, cyclohexane, benzene, toluene or xylene, or halogeno-hydrocarbons such as dichloromethane, dichloroethane, trichloromethane or chlorobenzene. It is also possible to use mixtures of the above-mentioned solvents. Preferred for the process is tetrahydrofuran or dioxan.

- Suitable acids for the process (II) + (III) + (IV) → (IB) are generally inorganic or organic acids or acid anhydrides. These preferably include carboxylic acids, such as, for example, acetic acid or
- 20 trifluoroacetic acid, sulfonic acids, such as, for example, methanesulfonic acid or p-toluenesulfonic acid, hydrochloric acid, phosphoric or phosphonic acids or anhydrides, such as polyphosphoric

acid or propanephosphonic acid anhydride. Preference is given to polyphosphoric acid ethyl ester. The acid is employed in an amount from 0.25 mol to 100 mol, relative to 1 mol of the compound of the general formula (III).

5 The process is in general carried out in a temperature range from +20°C to +150°C, preferably from +60°C to +100°C.

The process is generally carried out at normal pressure. However, it is also possible to carry it out at elevated pressure or at reduced pressure (for example in a range from 0.5 to 5 bar).

10 Suitable solvents for the process (IB) + (V) → (I) are generally customary organic solvents which do not change under the reaction conditions. These include ethers such as diethyl ether, diisopropyl ether, 1,2-dimethoxyethane, dioxan or tetrahydrofuran, ethyl acetate, acetone, acetonitrile, dimethylsulfoxide, dimethylformamide, or hydrocarbons such as pentane, hexane, cyclohexane, benzene, toluene or xylene, or halogeno-hydrocarbons such as dichloromethane, dichloroethane, trichloromethane or chlorobenzene. It is also possible to use mixtures of the above-mentioned solvents. Preferred for the process is tetrahydrofuran or dimethylformamide.

15 Suitable bases for the process (IB) + (V) → (I) are generally inorganic or organic bases. These preferably include alkali carbonates such as sodium or potassium carbonate or hydrogencarbonate, cyclic amines such as, for example, *N*-methylmorpholine, *N*-methylpiperidine, pyridine or 4-*N,N*-dimethylaminopyridine, or (C₁-C₄)-trialkylamines such as, for example, triethylamine or diisopropylethylamine, or alkali hydrides such as sodium or potassium hydride. Preference is given to
20 potassium carbonate or sodium hydride. The base is employed in an amount from 0.1 mol to 10 mol, preferably from 1 mol to 3 mol, relative to 1 mol of the compound of general formula (IV).

The process is in general carried out in a temperature range from 0°C to +150°C, preferably from +20°C to +80°C, especially at room temperature.

25 The process is generally carried out at normal pressure. However, it is also possible to carry it out at elevated pressure or at reduced pressure (for example in a range from 0.5 to 5 bar).


The compounds of the general formulas (II), (III), (IV) and (V) are known *per se*, or they can be prepared by customary methods.

30 The compounds of the present invention can also be prepared, if appropriate, by functional group transformations of individual substituents, especially those listed under R⁴ and R⁶, of the compounds of general formula (I) obtained by the process described above. These transformations are

carried out using standard synthetic methods, e.g. by esterification, ester cleavage / hydrolysis, amide formation, catalytic hydrogenation, alkylation and/or aryl coupling reactions.

The above-mentioned method can be illustrated by the following scheme:

Scheme



5

The compounds according to the invention exhibit an unforeseeable, useful pharmacological and pharmacokinetic activity spectrum. They are therefore suitable for use as medicaments for the treatment and/or prophylaxis of disorders in humans and animals.

Surprisingly, the compounds of the present invention show human neutrophil elastase (HNE) inhibitory activity and are therefore suitable for the preparation of medicaments for the treatment of diseases associated with HNE activity. They may thus provide an effective treatment of acute and chronic inflammatory processes, such as rheumatoid arthritis, atherosclerosis, and especially of acute and chronic pulmonary diseases, such as lung fibrosis, cystic fibrosis, pneumonia, acute

10

respiratory distress syndrome (ARDS), in particular pulmonary emphysema, including smoking-induced emphysema, and chronic obstructive pulmonary diseases (COPD), chronic bronchitis and bronchiectasis. The compounds of the present invention may further provide an effective treatment for cardiovascular ischaemic diseases such as acute coronary syndrome, acute myocardial infarction, unstable and stable angina pectoris, coronary artery bypass grafts (CABG) and heart failure development, for atherosclerosis, mitral valvular disease, atrial septal defects, percutaneous transluminal coronary angioplasty (PTCA), inflammation after open heart surgery and for pulmonary hypertension. They may also prove useful for an effective treatment of rheumatoid arthritis, acute inflammatory arthritis, cancer, acute pancreatitis, ulcerative colitis, periodontal disease, Chury-Strauss syndrome, acute and chronic atopic dermatitis, psoriasis, systemic lupus erythematosus, bullous pemphigus, sepsis, alcoholic hepatitis, liver fibrosis, Behcet's disease, allergic fungal sinusitis, allergic sinusitis, Crohn's disease, Kawasaki disease, glomerulonephritis, acute pyelonephritis, colorectal diseases, chronic suppurative otitis media, chronic venous leg ulcers, inflammatory bowel disease, bacterial and viral infections, brain trauma, stroke and other conditions in which neutrophil participation is involved.

The present invention further provides medicaments containing at least one compound according to the invention, preferably together with one or more pharmacologically safe excipient or carrier substances, and also their use for the above-mentioned purposes.

The active component can act systemically and/or locally. For this purpose, it can be applied in a suitable manner, for example orally, parenterally, pulmonally, nasally, sublingually, lingually, buccally, rectally, transdermally, conjunctivally, otically or as an implant.

For these application routes, the active component can be administered in suitable application forms.

Useful oral application forms include application forms which release the active component rapidly and/or in modified form, such as for example tablets (non-coated and coated tablets, for example with an enteric coating), capsules, sugar-coated tablets, granules, pellets, powders, emulsions, suspensions, solutions and aerosols.

Parenteral application can be carried out with avoidance of an absorption step (intravenously, intraarterially, intracardially, intraspinally or intralumbally) or with inclusion of an absorption (intramuscularly, subcutaneously, intracutaneously, percutaneously or intraperitoneally). Useful parenteral application forms include injection and infusion preparations in the form of solutions, suspensions, emulsions, lyophilisates and sterile powders.

Forms suitable for other application routes include for example inhalatory pharmaceutical forms (including powder inhalers, nebulizers), nasal drops/solutions, sprays; tablets or capsules to be administered linguallly, sublingually or buccally, suppositories, ear and eye preparations, vaginal capsules, aqueous suspensions (lotions, shake mixtures), lipophilic suspensions, ointments, creams, milk, pastes, dusting powders or implants.

The active components can be converted into the recited application forms in a manner known per se. This is carried out using inert non-toxic, pharmaceutically suitable excipients. These include inter alia carriers (for example microcrystalline cellulose), solvents (for example liquid polyethylene glycols), emulsifiers (for example sodium dodecyl sulphate), dispersing agents (for example polyvinylpyrrolidone), synthetic and natural biopolymers (for example albumin), stabilizers (for example antioxidants such as ascorbic acid), colorants (for example inorganic pigments such as iron oxides) or taste and/or odor corrigents.

For human use, in the case of oral administration, it is recommendable to administer doses of from 0.001 to 50 mg/kg, preferably of 0.01 mg/kg to 20 mg/kg. In the case of parenteral administration, such as, for example, intravenously or via mucous membranes nasally, buccally or inhalationally, it is recommendable to use doses of 0.001 mg/kg to 0.5 mg/kg.

In spite of this, it can be necessary in certain circumstances to depart from the amounts mentioned, namely as a function of body weight, application route, individual behaviour towards the active component, manner of preparation and time or interval at which application takes place. It can for instance be sufficient in some cases to use less than the aforementioned minimum amount, while in other cases the upper limit mentioned will have to be exceeded. In the case of the application of larger amounts, it can be advisable to divide them into a plurality of individual doses spread through the day.

The percentages in the tests and examples which follows are, unless otherwise stated, by weight; parts are by weight. Solvent ratios, dilution ratios and concentrations reported for liquid/liquid solutions are each based on the volume.

A. Evaluation of physiological activity

The potential of the compounds of the invention to inhibit neutrophil elastase activity may be demonstrated, for example, using the following assays:

I. In vitro enzyme assays of human neutrophil elastase (HNE)**5 Assay contents:**

assay buffer: 0.1 M HEPES-NaOH buffer pH 7.4, 0.5 M NaCl, 0.1% (w/v) bovine serum albumin;

suitable concentration (see below) of HNE (18 U/mg lyophil., #20927.01, SERVA Electrophoresis GmbH, Heidelberg, Germany) in assay buffer;

suitable concentration (see below) of substrate in assay buffer;

10 suitable concentration of test compounds diluted with assay buffer from a 10 mM stock solution in DMSO.

*Example I-A****In vitro* inhibition of HNE using a fluorogenic peptide substrate (continuous read-out signal, 384 MTP assay format):**

15 In this protocol, the elastase substrate MeOSuc-Ala-Ala-Pro-Val-AMC (#324740, Calbiochem-Novabiochem Corporation, Merck KGaA, Darmstadt, Germany) is used. The test solution is prepared by mixing 10 μ l of test compound dilution, 20 μ l of HNE enzyme dilution (final concentration 8 - 0.4 μ U/ml, routinely 2.1 μ U/ml) and 20 μ l of substrate dilution (final concentration 1 mM - 1 μ M, routinely 20 μ M), respectively. The solution is incubated for 0 - 2 hrs at
20 37°C (routinely one hour). The fluorescence of the liberated AMC due to the enzymatic reaction is measured at 37°C (TECAN spectra fluor plus plate reader). The rate of increase of the fluorescence (ex. 395 nm, em. 460 nm) is proportional to elastase activity. IC₅₀ values are determined by RFU-versus-[I] plots. K_m and K_{m(app.)} values are determined by Lineweaver-Burk plots and converted to K_i values by Dixon plots.

25 The preparation examples have IC₅₀ values within the range of 5 nM - 5 μ M in this assay. Representative data are given in Table 1:

Table 1

Example No.	IC ₅₀ [nM]
1	20
5	5
6	200
14	1000
21	130
43	23
45	20
69	50
73	1000
78	50

*Example I-B****In vitro* inhibition of HNE using a fluorogenic, insoluble elastin substrate (discontinuous read-out signal, 96 MTP assay format):**

- 5 In this protocol the elastase substrate elastin-fluorescein (#100620, ICN Biomedicals GmbH, Eschwege, Germany) is used. The test solution is prepared by mixing 3 µl of test compound dilution, 77 µl of HNE enzyme dilution (final concentration 0.22 U/ml - 2.2 mU/ml, routinely 21.7 µU/ml) and 80 µl substrate suspension (final concentration 2 mg/ml). The suspension is incubated for 0 - 16 hrs at 37°C (routinely four hours) under slightly shaking conditions. To stop the
- 10 enzymatic reaction, 160 µl of 0.1 M acetic acid are added to the test solution (final concentration 50 mM). The polymeric elastin-fluorescein is pulled down by centrifugation (Eppendorf 5804 centrifuge, 3.000 rpm, 10 min). The supernatant is transferred into a new MTP and the fluorescence of the liberated peptide fluorescein due to the enzymatic reaction is measured (BMG Fluostar plate reader). The rate of fluorescence (ex. 490 nm, em. 520 nm) is proportional to
- 15 elastase activity. IC₅₀ values are determined by RFU-versus-[I] plots.

II. In vitro human neutrophil assays

Example II-A

***In vitro* PMN elastolysis assay:**

This assay is used to determine the elastolytic potential of human polymorphonuclear cells (PMNs) and assess the proportion of degradation due to neutrophil elastase [cf. Z.W. She et al., *Am. J. Respir. Cell. Mol. Biol.* **9**, 386-392 (1993)].

Tritiated elastin, in suspension, is coated on to a 96 well plate at 10 µg per well. Test and reference [ZD-0892 (*J. Med. Chem.* **40**, 1876-1885, 3173-3181 (1997), WO 95/21855) and α1 protease inhibitor (α1PI)] compounds are added to the wells at the appropriate concentrations. Human PMNs are separated from peripheral venous blood of healthy donors and resuspended in culture media. The neutrophils are added to the coated wells at concentrations ranging between 1×10^6 to 1×10^5 cells per well. Porcine pancreatic elastase (1.3 µM) is used as a positive control for the assay, and α1PI (1.2 µM) is used as the positive inhibitor of neutrophil elastase. The cellular control is PMNs without compound at each appropriate cell density. The cells plus compounds are incubated in a humidified incubator at 37°C for 4 hours. The plates are centrifuged to allow the harvest of cell supernatant only. The supernatant is transferred in 75 µl volumes to corresponding wells of a 96 well Lumaplate™ (solid scintillant containing plates). The plates are dried until no liquid is visible in the wells and read in a beta counter for 3 minutes per well.

Elastolysis of the ^3H -elastin results in an increase in counts in the supernatant. An inhibition of this elastolysis shows a decrease, from the cellular control, of tritium in the supernatant. α1PI gave $83.46 \pm 3.97\%$ (mean ± s.e.m.) inhibition at 1.2 µM ($n = 3$ different donors at 3.6×10^5 cells per well). IC₅₀ values were obtained for the reference compound ZD-0892 of 45.50 ± 7.75 nM (mean ± s.e.m.) ($n = 2$ different donors at 3.6×10^5 cells per well).

Given that ZD-0892 is a selective inhibitor of PMN elastase along with the data from α1PI inhibition, these results indicate that the majority of elastin degradation by PMNs is due to the release of neutrophil elastase, and not to another elastolytic enzyme such as matrix metallo-proteases (MMPs). The compounds of this invention are evaluated for their inhibitory activity in this HNE-dependent model of neutrophil elastolysis.

*Example II-B****In vitro* inhibition of membrane bound elastase:**

Measurement of the inhibition of elastase bound to neutrophil membranes is performed using a human neutrophil assay. Neutrophils are stimulated with LPS at 37°C for 35 min and then spun at
5 1600 rpm. Subsequently, the membrane bound elastase is fixed to the neutrophils with 3% paraformaldehyde and 0.25% glutaraldehyde for 3 min at 4°C. The neutrophils are then spun, and vehicle and the compound under evaluation are added, followed by addition of the substrate MeOSuc-Ala-Ala-Pro-Val-AMC (#324740, Calbiochem-Novabiochem Corporation, Merck KGaA, Darmstadt, Germany) at 200 µM. Following a 25 min incubation at 37°C, the reaction is
10 terminated with PMSF (phenylmethanesulfonyl fluoride), and the fluorescence is read at ex: 400 nm and em: 505 nm. IC₅₀ values are determined by interpolation from plots of relative fluorescence vs. inhibitor concentration.

III. *In vivo* models*Example III-A***15 *In vivo* model of acute lung injury in the rat:**

Instillation of human neutrophil elastase (HNE) into rat lung causes acute lung damage. The extent of this injury can be assessed by measuring lung haemorrhage.

Rats are anaesthetised with Hypnorm/Hypnovel/water and instilled with HNE or saline delivered by microsyringe into the lungs. Test compounds are administered by intravenous injection, by oral
20 gavage or by inhalation at set times prior to the administration of HNE. Sixty minutes after the administration of elastase animals are killed by an anaesthetic overdose (sodium pentobarbitone) and the lungs lavaged with 2 ml heparinised phosphate buffered saline (PBS). Bronchoalveolar lavage (BAL) volume is recorded and the samples kept on ice. Each BAL sample is centrifuged at 900 r.p.m. for 10 minutes at 4-10°C. The supernatant is discarded and the cell pellet resuspended in
25 PBS and the sample spun down again. The supernatant is again discarded and the cell pellet resuspended in 1 ml 0.1% cetyltrimethyl-ammonium bromide (CTAB) / PBS to lyse the cells. Samples are frozen until blood content is assayed. Prior to the haemorrhage assay the samples are defrosted and mixed. 100 µl of each sample are placed into a separate well of a 96 well flat-bottomed plate. All samples are tested in duplicate. 100 µl 0.1% CTAB/PBS is included as a
30 blank. The absorbance of the well contents is measured at 415 nm using a spectrophotometer. A standard curve is constructed by measuring the OD at 415 nm of different concentrations of blood

in 0.1% CTAB/PBS. Blood content values are calculated by comparison to the standard curve (included in each plate) and normalised for the volume of BAL fluid retrieved.

The compounds of this invention are evaluated intravenously, orally or by inhalation for their inhibitory activity in this model of HNE-induced haemorrhage in the rat.

5 *Example III-B*

***In vivo* model of acute myocardial infarction in the rat:**

Elastase inhibitors are tested in a rat thread infarct model. Male Wistar rats (weighing >300 g) receive 10 mg/kg aspirin 30 min prior to surgery. They are anaesthetized by isofluran and ventilated (120-130 strokes/min, 200-250 µl stroke volume; MiniVent Type 845, Hugo Sachs
10 Elektronik, Germany) during the whole surgery. Following a left thoracotomy at the fourth inter-costal space, the pericardium is opened and the heart briefly exteriorized. A thread is turned around the left coronary artery (LAD) without occluding the artery. The thread is passed under the skin to the neck of the animal. The thorax is closed and the animal is allowed to recover for 4 days. At the fifth day, rats are anaesthetized with ether for 3 min, and the thread is tied and the LAD
15 occluded under ECG control. Test compounds are administered before or after LAD occlusion per os, intraperitoneally or intravenously (bolus or permanent infusion). After 1 hr occlusion, the thread is reopened to allow reperfusion. Hearts are excised, and infarct sizes are determined 48 hours later by staining of the re-occluded hearts with Evans blue, followed by TTC (triphenyltetrazolium chloride) staining of 2 mm heart sections. Normoxic (not occluded tissue)
20 areas stain blue, ischemic (occluded but surviving tissue) areas stain red and necrotic (occluded dead tissue) areas remain white. Each tissue section is scanned and infarct sizes are determined by computer planimetry.

B. Examples*Abbreviations:*

aq.	aqueous
c	concentration
conc.	concentrated
DMF	<i>N,N</i> -dimethylformamide
DMSO	dimethylsulfoxide
EI	electron impact ionisation (for MS)
ESI	electro-spray ionisation (for MS)
h	hour(s)
HPLC	high pressure liquid chromatography
LC-MS	liquid chromatography coupled with mass spectroscopy
min	minute(s)
Mp.	melting point
MS	mass spectroscopy
NMR	nuclear magnetic resonance spectroscopy
of th.	of theoretical (yield)
RP	reverse phase (for HPLC)
R _t	retention time (for HPLC)
THF	tetrahydrofuran

General methods:

5 All reactions are carried out under an argon atmosphere unless otherwise noted. Solvents are used as purchased from Aldrich without further purification. 'Silica gel' or 'Silica' refers to Silica gel 60 (0.040 mm-0.063 mm) from Merck KGaA company. Melting points are obtained with a Büchi 512 or similar melting point device and are uncorrected.

Compounds purified by preparative HPLC are purified over a RP18-column with acetonitrile and water as the eluent, using a 1:9 to 9:1 gradient.

*LC-MS / HPLC methods:****HPLC method 1***

Instrument: HP 1100 with DAD detection; column: Kromasil RP-18, 60 mm x 2 mm, 3.5 µm; eluent A: 5 ml HClO₄/l water, eluent B: acetonitrile; gradient: 0 min 2% B → 0.5 min 2% B → 4.5 min 90% B → 9 min 90% B; flow: 0.75 ml/min; oven: 30°C; UV detection: 210 nm.

LC-MS method 2

Instrument: Micromass Quattro LCZ with HPLC Agilent Series 1100; column: Grom-Sil120 ODS-4 HE, 50 mm x 2.0 mm, 3 µm; eluent A: 1 l water + 1 ml 50% formic acid, eluent B: 1 l acetonitrile + 1 ml 50% formic acid; gradient: 0.0 min 100% A → 0.2 min 100% A → 2.9 min 30% A → 3.1 min 10% A → 4.5 min 10% A; oven: 55°C; flow: 0.8 ml/min; UV detection: 208-400 nm.

LC-MS method 3

Instrument MS: Micromass ZQ; Instrument HPLC: Waters Alliance 2795; column: Phenomenex Synergi 2µ Hydro-RP Mercury 20 mm x 4 mm; eluent A: 1 l water + 0.5 ml 50% formic acid, eluent B: 1 l acetonitrile + 0.5 ml 50% formic acid; gradient: 0.0 min 90% A → 2.5 min 30% A → 3.0 min 5% A → 4.5 min 5% A; flow: 0.0 min 1 ml/min → 2.5 min/3.0 min/4.5 min 2 ml/min; oven: 50°C; UV detection: 210 nm.

HPLC method 4

Instrument MS: Micromass ZQ; Instrument HPLC: HP 1100 Series with DAD detection; column: Phenomenex Synergi 2µ Hydro-RP Mercury 20 mm x 4 mm; eluent A: 1 l water + 0.5 ml 50% formic acid, eluent B: 1 l acetonitrile + 0.5 ml 50% formic acid; gradient: 0.0 min 90% A → 2.5 min 30% A → 3.0 min 5% A → 4.5 min 5% A; flow: 0.0 min 1 ml/min → 2.5 min/3.0 min/4.5 min 2 ml/min; oven: 50°C; UV detection: 210 nm.

LC-MS method 5

Instrument: Micromass Quattro LCZ with HPLC Agilent Series 1100; column: Phenomenex Synergi 2µ Hydro-RP Mercury 20 mm x 4 mm; eluent A: 1 l water + 0.5 ml 50% formic acid, eluent B: 1 l acetonitrile + 0.5 ml 50% formic acid; gradient: 0.0 min 90% A → 2.5 min 30% A →

3.0 min 5% A → 4.5 min 5% A; flow: 0.0 min 1 ml/min → 2.5 min/3.0 min/4.5 min 2 ml/min; oven: 50°C; UV detection: 208-400 nm.


HPLC method 6

Instrument: HP 1100 with DAD detection; column: Kromasil RP-18, 60 mm x 2 mm, 3.5 µm;
5 eluent A: 5 ml HClO₄/l water, eluent B: acetonitrile; gradient: 0 min 2% B → 0.5 min 2% B → 4.5 min 90% B → 6.5 min 90% B; flow: 0.75 ml/min; oven: 30°C; UV detection: 210 nm.

Starting Materials and Intermediates:

Example 1A

Ethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-
10 pyrimidinecarboxylate




7.0 g (34.29 mmol) *N*-[3-(trifluoromethyl)phenyl]urea, 8.99 g (68.58 mmol) 4-cyanobenzaldehyde, 8.92 g (68.58 mmol) ethyl 3-oxobutanoate and 20 g polyphosphoric acid ethyl ester are suspended in 250 ml of tetrahydrofuran. The mixture is stirred at reflux for 18 hours. After cooling down to
15 room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.

Yield: 13.4 g (91% of th.)

¹H-NMR (200 MHz, DMSO-d₆): δ = 1.1 (t, 3H), 2.0 (s, 3H), 4.0 (q, 2H), 5.4 (d, 1H), 7.6 (m, 3H), 7.7 (m, 3H), 7.9 (m, 2H), 8.4 (d, 1H) ppm.

Example 2A

Allyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-pyrimidine-5-carboxylate




5 45.0 g ethyl polyphosphate are dissolved in 150 ml dioxane, 15.0 g (73.5 mmol) *N*-[3-(trifluoromethyl)phenyl]urea, 19.3 g (147 mmol) 4-cyanobenzaldehyde and 20.9 g (147 mmol) allyl acetoacetate are added and the mixture is stirred under reflux overnight. Volatiles are evaporated *in vacuo*, the remainder is dissolved in ethyl acetate and sequentially washed with saturated aqueous sodium hydrogencarbonate, sodium hydrogensulfite and sodium chloride
10 solution. The organic phase is dried over magnesium sulfate, filtered and evaporated to dryness *in vacuo*. The crude product is purified by column chromatography on silica gel (eluent: cyclohexane/ethyl acetate).

Yield: 18.4 g (50% of th.)

¹H-NMR (400 MHz, DMSO-d₆): δ = 2.08 (s, 3H), 4.55 (d, 2H), 5.05-5.18 (m, 2H), 5.41 (d, 1H),
15 5.82 (dddd, 1H), 7.54-7.92 (m, 8H), 8.41 (d, 1H) ppm.

Example 3A

Allyl (4*R*)-4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-pyrimidine-5-carboxylate




The enantiomers of Example 2A are separated by preparative HPLC on a chiral phase [chiral silica gel selector based on monomer *N*-methacryloyl-L-leucine-1-menthylamide, *cf.* EP-A-379 917; 250 mm x 20 mm; eluent: ethyl acetate → methanol → ethyl acetate; flow 50 ml/min; temperature
 5 24°C; detection 280 nm].

¹H-NMR (400 MHz, DMSO-d₆): δ = 2.08 (s, 3H), 4.55 (d, 2H), 5.05-5.18 (m, 2H), 5.41 (d, 1H), 5.82 (dddd, 1H), 7.54-7.92 (m, 8H), 8.41 (d, 1H) ppm.

[α]²⁰ = +25.9° (λ = 589 nm, methanol, c = 540 mg / 100 ml).

Example 4A

10 4-{5-Acetyl-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-4-pyrimidinyl}-benzonitrile




30 g (147 mmol) *N*-[3-(trifluoromethyl)phenyl]urea, 19.3 g (147 mmol) 4-cyanobenzaldehyde and 14.7 g (147 mmol) 2,4-pentanedione are suspended in 300 ml of tetrahydrofuran, and 90 g polyphosphoric acid ethyl ester are added. The mixture is stirred at reflux for 4 hours. After cooling down to room temperature, the solvent is removed *in vacuo*, the remainder is dissolved in ethyl acetate and sequentially washed with saturated aqueous sodium hydrogencarbonate and sodium chloride solution. The organic phase is dried over magnesium sulfate, filtered and evaporated to dryness *in vacuo*. The crude product is purified by column chromatography over silica gel (eluent: cyclohexane/ethyl acetate).

Yield: 16.8 g (29% of th.)

¹H-NMR (200 MHz, DMSO-d₆): δ = 2.0 (s, 3H), 2.2 (s, 3H), 5.5 (d, 1H), 7.5 (m, 1H), 7.6 (m, 3H), 7.7 (m, 1H), 7.8 (m, 1H), 7.9 (m, 2H), 8.5 (d, 1H) ppm.

Example 5A

4-(4-Cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidine-carboxylic acid



Method A:

3 g (7 mmol) of Example 1A are dissolved in a mixture of 50 ml water and 100 ml 5% potassium hydroxide in ethanol. The reaction mixture is stirred at room temperature for 18 hours. The solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with dichloromethane/methanol as eluent.

Yield: 1.27 g (45% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 2.0 (s, 3H), 5.4 (d, 1H), 7.6 (m, 1H), 7.6 (m, 2H), 7.7 (m, 1H), 7.8 (m, 1H), 7.9 (m, 3H), 8.3 (d, 1H), 12.5 (s, 1H) ppm.

Method B:


3.00 g (6.80 mmol) of Example 2A and 888 mg (10.2 mmol) morpholine are dissolved under argon
5 in 30 ml tetrahydrofuran at room temperature. 392 mg (0.34 mmol) tetrakis(triphenylphosphine)-
palladium(0) are added, and the mixture is reacted for 15 minutes at room temperature. The solvent
is evaporated *in vacuo*, the remainder is dissolved in ethyl acetate and washed sequentially with 2
N hydrochloric acid, water and saturated sodium chloride solution. The organic phase is dried over
magnesium sulfate and evaporated to dryness. The crude product is purified by preparative RP-
10 HPLC with a water/acetonitrile gradient.

Yield: 1.51 g (52% of th.)

¹H-NMR: see above.

Example 6A

(4*R*)-4-(4-Cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-
15 pyrimidinecarboxylic acid



Method A:

The enantiomers of Example 5A are separated by preparative HPLC on a chiral phase [chiral silica gel selector based on monomer *N*-methacryloyl-L-leucine-1-menthylamide, *cf.* EP-A-379 917;

250 mm x 20 mm; eluent: ethyl acetate → methanol → ethyl acetate; flow 25 ml/min; temperature 23°C; detection 254 nm].

¹H-NMR (300 MHz, DMSO-d₆): δ = 2.0 (s, 3H), 5.4 (d, 1H), 7.6 (m, 1H), 7.6 (m, 2H), 7.7 (m, 2H), 7.8 (m, 1H), 7.9 (m, 2H), 8.3 (d, 1H), 12.5 (s, 1H) ppm.


5 [α]²⁰ = +2.5° (λ = 589 nm, methanol, c = 505 mg / 100 ml).

Method B:

In analogy to Example 5A (method B), this compound is prepared from Example 3A in 87% yield.

Example 7A

2-Hydroxyethyl (4*R*)-4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-
10 tetrahydro-5-pyrimidinecarboxylate



Under argon, 1560 mg (3.89 mmol) of Example 6A are added to 19.6 ml dimethylformamide. After addition of 1.095 ml (7.86 mmol) triethylamine and 1.11 ml (15.7 mmol) 2-bromoethanol, the reaction mixture is stirred at ca. 70°C for 8 hours. After cooling to room temperature, the
15 reaction mixture is concentrated *in vacuo*. The residue is taken up in ethyl acetate and washed with water. After drying with magnesium sulfate, the organic phase is evaporated *in vacuo*. The residue is taken up in 8 ml methanol and purified by preparative HPLC (column: Nucleosil 100-5 C18 Nautilus, 20 x 50 mm, 5 μm; solvent A: acetonitrile, solvent B: water + 0.3% formic acid; gradient: 0 min 10% A → 2 min 10% A → 6 min 90% A → 7 min 90% A → 7.1 min 10% A → 8
20 min 10% A; wavelength: 220 nm; injection volume: ca. 500 μl; number of injections: 18). The product containing fractions are combined and lyophilized.

Yield: 1290 mg (74.5% of th.)


MS (EI): $m/z = 446$ ($M+H$)⁺

¹H-NMR (300 MHz, DMSO-d₆): $\delta = 2.05$ (d, 3H), 3.5 (q, 2H), 3.95-4.15 (m, 2H), 4.75 (tr, 1H), 5.45 (d, 1H), 7.55-7.75 (m, 5H), 7.75 (d, 1H), 7.85 (d, 2H), 8.35 (d, 1H) ppm.

5 $[\alpha]^{20} = +14.3^\circ$ ($\lambda = 589$ nm, methanol, $c = 455$ mg / 100 ml).

Example 8A

5-(Benzyloxy)-hexane-2,4-dione



To a solution of ethyl 3-(benzyloxy)-2-methylpropanoate (13.4 g, 64 mmol) in dimethylsulfoxide
10 (50 ml) is added sodium hydride (2.57 g, 64.34 mmol; 60% dispersion in mineral oil). After 5
minutes stirring, a solution of acetone (2.37 ml, 32.1 mmol) in dimethylsulfoxide (30 ml) is added,
and the reaction is stirred at 60°C overnight. After cooling to room temperature, saturated aqueous
ammonium chloride (100 ml) is added and the product is extracted with ethyl acetate (3 x 150 ml).
The combined organic phases are washed with brine, dried over anhydrous magnesium sulfate,
15 filtered and concentrated. The residue is purified by flash chromatography over silica gel 60 (300
g) using cyclohexane / ethyl acetate (10:1 to 5:1) as eluent. The title compound is isolated as a
mixture of enol ethers.


Yield: 4.92 g (69% of th.)

HPLC (method 1): $R_t = 4.17$ min and 4.75 min, $\lambda_{\max} = 280$ nm

20 MS (ESIpos): $m/z = 220$ ($M+H$)⁺

Example 9A

1-Benzyl 2-ethyl piperidine-1,2-dicarboxylate



To a stirred solution of ethyl piperidine-2-carboxylate (15 g, 95 mmol), triethylamine (27 ml, 191 mmol) and 4-dimethylaminopyridine (0.58 g, 4.8 mmol) in dichloromethane (100 ml) at 0°C is added a solution of benzyl chloridocarbonate (17 g, 100 mmol) dropwise. The reaction is allowed to warm slowly to room temperature. It is stirred at room temperature overnight (16 h), then allowed to stand for 2 days. The crude product is extracted with dichloromethane, washed with 1 N hydrochloric acid, saturated aqueous sodium bicarbonate solution and brine. The organic phase is dried with anhydrous magnesium sulfate, filtered and concentrated *in vacuo*. The residue is purified by flash chromatography over silica gel 60 (300 g) with cyclohexane / ethyl acetate as eluent to afford a pale yellow oil.


Yield: 17.9 g (62% of th.)

HPLC (method 6): $R_t = 4.91$ min, $\lambda_{\max} = 202$ nm

MS (ESIpos): $m/z = 309$ ($M + NH_4$)⁺

Example 10A

15 Benzyl 2-acetoacetylpiperidine-1-carboxylate



The title compound is prepared according to the procedure described in Example 8A and is isolated as the enol ether.

Yield: 2.13 g (82% of th.)


HPLC (method 1): $R_t = 4.45$ and 4.98 min, $\lambda_{\max} = 276$ nm

5 MS (ESIpos): $m/z = 321(M+NH_4)^+$

1H -NMR (300 MHz, $CDCl_3$): $\delta = 15.4$ (br s, 1H), 7.45-7.20 (m, 5H), 5.62-4.68 (m, 3H), 4.29-3.63 (m, 2H), 3.15-2.80 (m, 1H), 2.37-1.10 (m, 6H), 2.01 (s, 3H) ppm.

Example 11A

2-(Benzyloxy)-1-[(benzyloxy)methyl]ethyl 3-oxobutanoate



10


1.00 g (3.67 mmol) 1,3-dibenzyloxy-2-propanol is dissolved in 10 ml toluene, 3.72 mg (0.04 mmol) triethylamine are added, and the mixture is stirred at 90°C. 401 mg (4.77 mmol) diketene are added, and stirring is continued for 1 h. The mixture is cooled to room temperature, diluted with ice-water and extracted with toluene. The organic layer is dried over magnesium sulfate and
15 evaporated to dryness. The crude product is purified by column chromatography on silica gel (eluent: cyclohexane/ethylacetate 2:1).

Yield: 1.03 g (69% of th.)

1H -NMR (300 MHz, $DMSO-d_6$): $\delta = 2.15$ (s, 3H), 3.55-3.65 (m, 6H), 4.44 (dd, 2H), 5.67 (q, 1H), 7.23-7.40 (m, 10H) ppm.

Example 12A

(1*S*)-2-(Benzyloxy)-1-methyl-2-oxoethyl 3-oxobutanoate




In analogy to the procedure of Example 11A, the title compound is synthesized using (*S*)-benzyl
5 lactate with a yield of 80% of th.

¹H-NMR (200 MHz, DMSO-d₆): δ = 1.42 (d, 3H), 2.18 (s, 3H), 3.66 (s, 2H), 5.11 (q, 1H), 5.17 (s, 2H), 7.30-7.43 (m, 5H) ppm.

Example 13A

Allyl 3-(2-*tert.*-butoxy-2-oxoethyl)-4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)-
10 phenyl]-1,2,3,4-tetrahydropyrimidine-5-carboxylate




1000 mg (2.27 mmol) of Example 2A are dissolved in 10 ml dimethylformamide, 344 mg (2.49 mmol) potassium carbonate and 486 mg (2.49 mmol) *tert.*-butyl bromoacetate are added, and the suspension is stirred at room temperature overnight. The mixture is partitioned between ethyl
15 acetate and aqueous potassium dihydrogenphosphate/disodium hydrogenphosphate buffer (pH 7). The combined organic extracts are washed with water and aqueous sodium chloride solution, dried over magnesium sulfate, and evaporated *in vacuo*. The crude product is purified by column chromatography on silica gel (eluent: cyclohexane/ethyl acetate 3:1).

Yield: 985 mg (78% of th.)

$^1\text{H-NMR}$ (200 MHz, DMSO-d_6): δ = 1.29 (s, 9H), 2.08 (s, 3H), 3.88 (d, 1H), 4.09 (d, 1H), 4.52 (d, 2H), 5.09-5.15 (m, 2H), 5.60 (s, 1H), 5.71-5.92 (m, 1H), 7.60-7.93 (m, 8H) ppm.

Example 14A

- 5 3-(2-*tert.*-Butoxy-2-oxoethyl)-4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidine-5-carboxylic acid




- 985 mg (1.77 mmol) of Example 13A and 231 mg (2.66 mmol) morpholine are dissolved in 10 ml tetrahydrofuran at room temperature under an argon atmosphere. 102 mg (0.09 mmol) tetrakis-
10 (triphenylphosphine)palladium(0) are added, and the solution is stirred for 30 minutes. The solvent is evaporated *in vacuo*, the remainder is dissolved in ethyl acetate and washed with 2 M hydrochloric acid, water and aqueous sodium chloride solution, dried over magnesium sulfate and evaporated *in vacuo*. The crude product is purified using RP-HPLC with a water/acetonitrile gradient.

- 15 Yield: 662 mg (70% of th.)

$^1\text{H-NMR}$ (200 MHz, DMSO-d_6): δ = 1.30 (s, 9H), 2.05 (s, 3H), 3.85 (d, 1H), 4.09 (d, 1H), 5.55 (s, 1H), 7.57-7.90 (m, 8H), 12.6 (br s, 1H) ppm.

Example 15A

- tert.*-Butyl [6-(4-cyanophenyl)-5-(1*H*-imidazol-1-ylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoro-
20 methyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate




372 mg (0.72 mmol) of Example 14A are dissolved in 5 ml dimethylformamide, 351 mg (2.17 mmol) 1,1'-carbonyldiimidazole are added, and the mixture is stirred at room temperature overnight. The mixture is partitioned between water and ethyl acetate, the combined organic extracts
 5 are washed with aqueous sodium hydrogencarbonate solution, water and aqueous sodium chloride solution, dried over magnesium sulfate, filtered and evaporated to dryness *in vacuo*.

Yield: 392 mg (90% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 1.30 (s, 9H), 1.48 (s, 3H), 3.83 (d, 1H), 4.05 (d, 1H), 5.65 (s, 1H), 7.05 (s, 1H), 7.62 (s, 1H), 7.68-7.90 (m, 7H), 7.98 (s, 1H), 8.37 (s, 1H) ppm.

10 Example 16A

4-{5-[(1H-1,2,3-Benzotriazol-1-yloxy)carbonyl]-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl}benzonitrile




500 mg (1.25 mmol) of Example 5A, 310 mg (1.62 mmol) 1-ethyl-3-(3-(dimethylamino)propyl)-carbodiimid-hydrochloride and 202 mg (1.49 mmol) 1-hydroxybenzotriazole are dissolved in 2 ml dimethylformamide and stirred at room temperature overnight. The reaction mixture is partitioned between ethyl acetate and water, the organic phase is washed with saturated sodium chloride solution, dried over magnesium sulfate and evaporated to dryness *in vacuo*. The crude product is used directly for further reactions.

Yield: 580 mg (76% of th.)

¹H-NMR (400 MHz, DMSO-d₆): δ = 2.17 (s, 3H), 5.82 (d, 1H), 7.23 (d, 1H), 7.35-8.03 (m, 10H), 8.10 (d, 1H), 8.82 (d, 1H) ppm.

10 Example 17A

Tribenzyl 2-{4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl}ethane-1,1,1-tricarboxylate




200 mg (0.52 mmol) of Example 1, 259 mg (0.62 mmol) tribenzyl methanetricarboxylate and 203 mg (0.77 mmol) triphenylphosphine are dissolved in 3 ml tetrahydrofuran under an argon atmosphere. The solution is cooled to 0°C, and 156 mg (0.77 mmol) diisopropyl azodicarboxylate are added slowly. The mixture is warmed to room temperature overnight, evaporated to dryness and purified directly by vacuum flash chromatography on silica (eluent: petrol ether/ethyl acetate 2:1 → 1:1) and thereafter by RP-HPLC (eluent: acetonitrile/water gradient).

Yield: 60 mg (14% of th.)

¹H-NMR (400 MHz, DMSO-d₆): δ = 1.32 (s, 3H), 2.74 (d, 1H), 3.21 (d, 1H), 4.80 (d, 1H), 5.16 (s, 6H), 7.18-7.33 (m, 15H), 7.38 (d, 1H), 7.46 (s, 1H), 7.54 (d, 2H), 7.62 (t, 1H), 7.70 (d, 1H), 7.32-7.38 (m, 3H) ppm.

Example 18A

- 5 (4*R*)-4-{5-Acetyl-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-4-pyrimidinyl}-benzonitrile




- The enantiomers of Example 4A are separated by preparative HPLC on a chiral phase [chiral silica gel selector based on monomer *N*-methacryloyl-L-leucine-1-menthylamide, *cf.* EP-A-379 917; 10 250 mm x 20 mm; eluent: ethyl acetate → methanol → ethyl acetate; flow 25 ml/min; temperature 23°C; detection 254 nm].

¹H-NMR (200 MHz, DMSO-d₆): δ = 2.0 (s, 3H), 2.2 (s, 3H), 5.5 (d, 1H), 7.5 (m, 1H), 7.6 (m, 3H), 7.7 (m, 1H), 7.8 (m, 1H), 7.9 (m, 2H), 8.5 (d, 1H) ppm.

[α]²⁰ = +45.9° (λ = 589 nm, methanol, c = 530 mg / 100 ml).

- 15 *Example 19A*

(4*R*)-Tribenzyl 2-{4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl}ethane-1,1,1-tricarboxylate




1.0 g (2.58 mmol) of Example 4, 1.3 g (3.10 mmol) tribenzyl methanetricarboxylate and 1.02 g (3.87 mmol) triphenylphosphine are dissolved in 15 ml tetrahydrofuran under an argon atmosphere. The solution is cooled to 0°C, and 0.67 g (3.87 mmol) diisopropyl azodicarboxylate are added slowly. The mixture is warmed to room temperature overnight, evaporated to dryness and purified directly by vacuum flash chromatography on silica (eluent: cyclohexane/ethyl acetate 2:1 → 1:1).

Yield: 0.65 mg (31% of th.)

¹H-NMR (400 MHz, DMSO-d₆): δ = 1.3 (s, 3H), 2.7 (d, 1H), 3.2 (d, 1H), 4.8 (d, 1H), 5.2 (s, 6H), 7.2 (m, 6H), 7.3 (m, 9H), 7.4 (m, 1H), 7.5 (m, 1H), 7.5 (m, 2H), 7.6 (m, 1H), 7.7 (m, 1H), 7.8 (m, 3H) ppm.

Example 20A

1-(2,2-Dimethyl-1,3-dioxolan-4-yl)-butane-1,3-dione



To sodium hydride (3.7 g, 93 mmol; 60% dispersion in mineral oil) under an argon atmosphere is added DMSO (30 ml) followed by a solution of methyl 2,2-dimethyl-1,3-dioxolane-4-carboxylate (15 g, 93 mmol) in DMSO (50 ml). A solution of acetone (2.7 g, 45 mmol) in DMSO (50 ml) is then added dropwise over 1 hour. The solution is stirred at room temperature for 16 h, then


quenched with saturated aqueous ammonium chloride solution (350 ml) and extracted with diethyl ether (1 l). The ether layer is washed with brine, dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo*. The crude product is chromatographed over silica gel 60 with dichloromethane as eluent. The compound is isolated as its enol ether.

5 Yield: 3.98 g (46% of Th.)

¹H-NMR (300 MHz, CDCl₃): δ = 5.90 (s, 1H), 4.56-4.44 (m, 1H), 4.32-4.23 (m, 1H), 4.04-3.96 (m, 1H), 2.11 (s, 3H), 1.49 (s, 3H), 1.41 (s, 3H) ppm.

*Preparation Examples:**Example 1*

4-{5-(Hydroxymethyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl} benzonitrile



5

2.05 g (4.77 mmol) of Example 1A are dissolved in 40 ml tetrahydrofuran. At 0°C, 9.55 ml (9.55 mmol) of 1 M lithium aluminiumhydride in tetrahydrofuran are added dropwise. The solution is stirred at 0°C for two hours and then quenched with methanol. The solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with dichloromethane/methanol mixtures as eluent.


10

Yield: 1.08 g (58% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 1.5 (s, 3H), 3.7 (dd, 1H), 4.1 (dd, 1H), 4.8 (dd, 1H), 5.1 (d, 1H), 7.5-7.7 (m, 6H), 7.9 (m, 3H) ppm.

Examples 2 and 3

15 4-{5-(1-Hydroxyethyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl} benzonitrile



200 mg (0.5 mmol) of Example 4A and 49 mg (0.6 mmol) sodium acetate are dissolved in 5 ml ethanol. 47 mg (1.25 mmol) sodium borohydride are added. The solution is stirred at room temperature for 16 hours, then water is added. The product precipitates as mixture of diastereomers (94 mg). The diastereomers are separated by preparative HPLC.

Example 2 (Diastereomer I):

Yield: 52 mg (26% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 0.8 (d, 3H), 1.6 (s, 3H), 4.6 (m, 1H), 4.9 (d, 1H), 5.1 (d, 1H), 7.5-7.7 (m, 6H), 7.8 (d, 1H), 7.9 (m, 2H) ppm.


Example 3 (Diastereomer II):

Yield: 41 mg (20% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 1.2 (d, 3H), 1.6 (s, 3H), 4.6 (m, 2H), 4.9 (d, 1H), 7.5 (m, 2H), 7.6 (m, 4H), 7.8 (m, 2H), 7.9 (d, 1H) ppm.

Example 4

(4*R*)-4-{5-(Hydroxymethyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl} benzonitrile




3.00 g (6.80 mmol) of Example 3A are dissolved in 20 ml dry tetrahydrofuran, 6.80 ml (6.80 mmol) of 1 M lithium aluminiumhydride in tetrahydrofuran are added slowly, and stirring is continued for 30 minutes at 0°C. Saturated ammonium chloride solution is added cautiously to hydrolyze excess of lithium aluminiumhydride, followed by 50 ml ethyl acetate and dropwise addition of water to precipitate inorganic salts. The organic phase is decanted, washed with water, dried over magnesium sulfate and evaporated to dryness. The crude product is purified by column chromatography over silica gel (eluent: dichloromethane/methanol 100:0 → 95:5).

Yield: 1.38 g (40% of th.)

¹H-NMR (200 MHz, DMSO-d₆): δ = 1.55 (s, 3H), 3.67 (dd, 1H), 4.10 (dd, 1H), 4.77 (t, 1H), 5.09 (d, 1H), 7.50 (d, 1H), 7.55-7.75 (m, 5H), 7.82 (d, 1H), 7.87 (d, 2H) ppm.

Example 5

(4R)-[5-{[2-(Carboxymethoxy)ethoxy]carbonyl}-6-(4-cyanophenyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2H)-yl]acetic acid




80 mg (0.18 mmol) of Example 7A are dissolved in 4 ml tetrahydrofuran, and 36 mg (0.9 mmol) sodium hydride (60% suspension in mineral oil) are added. After stirring for one hour at room temperature, 50 mg (0.36 mmol) bromoacetic acid are added. After stirring at room temperature for
 5 four hours, the mixture is quenched with methanol, the solvent is removed *in vacuo* and the residue is purified by preparative HPLC.

Yield: 21 mg (21% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 2.0 (s, 3H), 2.9 (d, 1H), 3.5-3.7 (m, 3H), 4.0 (m, 2H), 4.3 (m, 2H), 5.9 (s, 1H), 7.6-7.7 (m, 6H), 7.8 (m, 3H) ppm.

10 Example 6

4-(4-Cyanophenyl)-*N*-cyclopropyl-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidine-5-carboxamide





100 mg (0.25 mmol) of Example 5A are dissolved in 2 ml tetrahydrofuran, and 3 mg (0.02 mmol) 4-*N,N*-dimethylaminopyridine, 39 mg (0.3 mmol) *N,N*-diisopropylethylamine and 156 mg (0.3 mmol) benzotriazol-1-yloxy-tris(pyrrolidino)phosphonium hexafluorophosphate are added. The reaction mixture is stirred at room temperature for 15 minutes, then 29 mg (0.5 mmol) cyclopropylamine are added. The reaction mixture is stirred at room temperature for 72 hours. The mixture is evaporated to dryness *in vacuo* and the crude product is purified by preparative HPLC.

Yield: 71 mg (62% of th.)

LC-MS (method 2): $R_t = 3.98$ min


MS (ESIpos): $m/z = 441$ ($M+H$)⁺.

10 In analogy to the procedure for Example 6, the following compounds are prepared:

Example No.	Structure	Starting materials	Yield [%]	R_t [min] (method)	Mass [$M+H$] ⁺
7		Example 5A; 1-(3,4-dichloro-phenyl)-piperazine	61	4.32 (2)	614
8		Example 5A; 1-benzyl-piperidin-4-amine	26	3.49 (2)	574


Example No.	Structure	Starting materials	Yield [%]	R _t [min] (method)	Mass [M+H] ⁺
9		Example 5A; methyl piperazine-1- carboxylate	66	3.96 (2)	528
10		Example 5A; 1-pyridin-4-yl- piperazine	30	3.34 (2)	547
11		Example 5A; 1-(2-chloro- phenyl)methan- amine	44	4.20 (2)	525
12		Example 5A; 2-fluoroaniline	20	4.24 (2)	495

Example No.	Structure	Starting materials	Yield [%]	R _t [min] (method)	Mass [M+H] ⁺
13		Example 5A; 2-(2-fluoro-phenyl)ethan-amine	59	4.19 (2)	523
14		Example 5A; 1-(2-fluoro-phenyl)methan-amine	56	4.15 (2)	509
15		Example 5A; 1,4'-bipiperidine	23	3.35 (2)	552
16		Example 5A; 1-propyl-piperazine	64	3.24 (2)	514

Example No.	Structure	Starting materials	Yield [%]	R _t [min] (method)	Mass [M+H] ⁺
17		Example 5A; ethyl piperidine-4- carboxylate	73	4.10 (2)	541

Examples 18 and 19

4-{5-[2-(Benzyloxy)propanoyl]-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl}benzonitrile



5

To a stirred solution of 5-(benzyloxy)hexane-2,4-dione (250 mg, 1.13 mmol; Example 8A), 4-cyanobenzaldehyde (149 mg, 1.13 mmol) and *N*-[3-(trifluoromethyl)phenyl]urea (232 mg, 1.13 mmol) in tetrahydrofuran (5 ml) is added polyphosphoric acid ethyl ester (700 mg). After 16 hours, the reaction solution is purified directly by preparative HPLC (RP18 column; eluent: acetonitrile/0.1% aq. formic acid 10:90 → 90:10). The fractions containing product are concentrated and again purified by preparative HPLC (RP18 column; eluent: acetonitrile/0.1% aq. formic acid 10:90 → 90:10).

Example 18 (Diastereomer I):

Yield: 28 mg (4.75% of th.)

HPLC (method 1): $R_t = 4.86$ min, $\lambda_{\max} = 234$ nm

MS (ESIpos): $m/z = 520$ (M+H)⁺

- 5 ¹H-NMR (300 MHz, DMSO-d₆): $\delta = 8.46$ (d, 1H, $J = 2.8$ Hz), 7.93-7.12 (m, 13H), 5.59 (d, 1H, $J = 2.8$ Hz), 4.30-4.11 (m, 3H), 1.87 (s, 3H), 1.32-1.11 (m, 3H) ppm.

Example 19 (Diastereomer II):

Yield: 24.3 mg (4.12% of th.)


HPLC (method 1): $R_t = 4.93$ min, $\lambda_{\max} = 232$ nm

- 10 MS (ESIpos): $m/z = 520$ (M+H)⁺

¹H-NMR (300 MHz, DMSO-d₆): $\delta = 8.48$ (d, 1H, $J = 3.2$ Hz), 7.90-7.13 (m, 13H), 5.48 (d, 1H, $J = 2.8$ Hz), 4.50-4.27 (m, 3H), 1.88 (s, 3H), 1.14 (d, 3H, $J = 6.4$ Hz) ppm.

Example 20

- 15 4-{5-(Cyclobutylcarbonyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-pyrimidin-4-yl} benzonitrile



To a stirred suspension of 1-cyclobutylbutane-1,3-dione (69 mg, 0.49 mmol), *N*-[3-(trifluoromethyl)phenyl]urea (100 mg, 0.49 mmol) and 4-cyanobenzaldehyde (64.2 mg, 0.49 mmol) in

tetrahydrofuran (250 ml) is added polyphosphoric acid ethyl ester (300 mg). The reaction mixture is stirred at reflux for 18 hours. After cooling to room temperature, the solvent is removed *in vacuo* and the crude product is purified by preparative HPLC (RP18 column; eluent: acetonitrile/0.1% aq. formic acid 10:90 → 90:10).

5 Yield: 54.6 mg (20% of th.)

LC-MS (method 5): $R_t = 2.57$ min


HPLC (method 6): $R_t = 4.91$ min, $\lambda_{\max} = 200$ nm

MS (ESIpos): $m/z = 457$ ($M + \text{NH}_4$)⁺

¹H-NMR (300 MHz, DMSO- d_6): $\delta = 8.39$ (d, 1H, $J = 3.7$ Hz), 7.98-7.48 (m, 8H), 5.36 (d, 1H, $J =$
10 3.7 Hz), 3.56-3.43 (m, 1H), 2.22-1.40 (m, 6H), 1.94 (s, 3H) ppm.

Example 21

4-{5-(Cyclohexylcarbonyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-pyrimidin-4-yl} benzonitrile



15 The title compound is prepared according to the procedure described for Example 20.

Yield: 72.8 mg (27% of th.)

LC-MS (method 3): $R_t = 2.63$ min


HPLC (method 6): $R_t = 5.18$ min, $\lambda_{\max} = 196$ nm

MS (ESIpos): $m/z = 485$ ($M+NH_4$)⁺

¹H-NMR (300 MHz, DMSO-d₆): $\delta = 8.35$ (d, 1H), 7.95-7.47 (m, 8H), 5.47 (d, 1H), 2.69 (m, 1H), 1.84 (s, 3H), 1.88-0.92 (m, 10H) ppm.

Example 22

- 5 4-{5-(Cyclopropylcarbonyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl}benzonitrile



The title compound is prepared according to the procedure described for Example 18.

Yield: 113 mg (52% of th.)

- 10 LC-MS (method 5): $R_t = 2.46$ min


HPLC (method 6): $R_t = 4.72$ min, $\lambda_{\max} = 202$ nm

MS (ESIpos): $m/z = 443$ ($M+NH_4$)⁺

¹H-NMR (300 MHz, DMSO-d₆): $\delta = 8.38$ (d, 1H, $J = 3.6$ Hz), 7.91-7.51 (m, 8H), 5.53 (d, 1H, $J = 3.4$ Hz), 3.56-3.43 (m, 1H), 1.98 (s, 3H), 0.93-0.72 (m, 4H) ppm.

- 15 *Example 23*

(4*R*)-4-{5-(Cyclopropylcarbonyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl}benzonitrile




The enantiomers of Example 22 are separated by preparative HPLC on a chiral phase [chiral silica gel selector based on poly(*N*-methacryloyl-L-leucine-dicyclopropylmethylamide); column: 250 mm x 20 mm; gradient: 0-6 minutes ethyl acetate, 6-8 minutes methanol, 8-10 minutes ethyl acetate; flow: 25 ml/min; detection: UV 254 nm].

$[\alpha]^{20} = +32.0^\circ$ ($\lambda = 589$ nm, methanol, $c = 633$ mg / 100 ml).

Example 24

4-{5-(4-Methoxybenzoyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-pyrimidine-4-yl}benzonitrile



10

The title compound is prepared according to the procedure described for Example 18.

Yield: 69.7 mg (28% of th.)

LC-MS (method 4): $R_t = 2.58$ min


HPLC (method 6): $R_t = 4.83$ min, $\lambda_{\max} = 200$ nm

MS (ESIpos): $m/z = 492$ (M+H)⁺

¹H-NMR (300 MHz, CDCl₃): $\delta = 7.75$ -7.39 (m, 10H), 6.89 (d, 2H), 5.67 (m, 2H), 3.85 (s, 3H),
5 1.49 (s, 3H) ppm.

Example 25

4-{5-[(2-Amino-4-methyl-1,3-thiazol-5-yl)carbonyl]-6-methyl-2-oxo-1-[3-(trifluoromethyl)-phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl} benzonitrile



10 The title compound is prepared according to the procedure described for Example 18.

Yield: 63 mg (26% of th.)


LC-MS (method 3): $R_t = 2.22$ min

MS (ESIpos): $m/z = 498$ (M+H)⁺

¹H-NMR (400 MHz, CDCl₃): $\delta = 7.78$ -7.20 (m, 9H), 6.56-6.40 (m, 2H), 5.69 (br s, 1H), 2.24 (s,
15 3H), 1.93 (s, 3H) ppm.

Example 26

tert.-Butyl [6-(4-cyanophenyl)-5-(cyclobutylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)-phenyl]-3,6-dihydropyrimidine-1(2H)-yl]acetate



A stirred suspension of 4-{5-(cyclobutylcarbonyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl}benzonitrile (Example 20) (29 mg, 0.066 mmol) and potassium carbonate (16.4 mg, 0.12 mmol) in dimethylformamide (1 ml) is treated with *tert.*-butyl bromoacetate (14.2 mg, 0.073 mmol), and then stirred at room temperature overnight (16 h). The reaction solution is quenched with water (3 ml) and extracted with diethyl ether (2 x 5 ml). The combined organic phases are washed with brine, dried over anhydrous magnesium sulfate, filtered and concentrated.

Yield: 40 mg (94.4% of th.)

10 LC-MS (method 4): $R_t = 3.05$ min


HPLC (method 1): $R_t = 5.43$ min, $\lambda_{\max} = 234$ nm

MS (ESIpos): $m/z = 554$ ($M+H$)⁺

¹H-NMR (300 MHz, DMSO-*d*₆): $\delta = 7.80$ -7.55 (m, 8H), 5.58 (br s, 1H), 4.19-3.91 (m, 2H), 3.60-3.47 (m, 1H), 2.40-0.85 (m, 6H), 1.94 (s, 3H), 1.26 (s, 9H) ppm.

15 Example 27

tert.-Butyl [6-(4-cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate



The title compound is prepared from Example 22 according to the procedure described for Example 26.

Yield: 89 mg (100% of th.)

5 LC-MS (method 4): $R_t = 2.93$ min


HPLC (method 1): $R_t = 5.27$ min, $\lambda_{\max} = 200$ nm

MS (ESIpos): $m/z = 540$ ($M+H$)⁺

¹H-NMR (300 MHz, DMSO-d₆): $\delta = 7.95$ -7.54 (m, 8H), 5.73 (s, 1H), 4.16-3.83 (m, 2H), 2.23 (m, 1H), 1.98 (s, 3H), 1.29 (s, 9H), 0.91-0.72 (m, 4H) ppm.

10 Example 28

tert.-Butyl [6-(4-cyanophenyl)-5-(cyclohexylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)-phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate



The title compound is prepared from Example 21 according to the procedure described for Example 26.

Yield: 65 mg (90% of th.)


5 HPLC (method 1): $R_t = 5.65$ min, $\lambda_{\max} = 234$ nm

MS (ESIpos): $m/z = 582$ ($M+H$)⁺

¹H-NMR (300 MHz, DMSO-*d*₆): $\delta = 7.90$ - 7.52 (m, 8H), 5.68 (s, 1H), 4.20 - 3.88 (m, 2H), 2.78 - 2.66 (m, 1H), 1.72 (s, 3H), 1.69 - 1.00 (m, 10H), 1.29 (s, 9H) ppm.

Example 29

10 *tert*.-Butyl [5-[(2-amino-4-methyl-1,3-thiazol-5-yl)carbonyl]-6-(4-cyanophenyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate



The title compound is prepared from Example 25 according to the procedure described for Example 26.

Yield: 38 mg (60% of th.)


LC-MS (method 4): $R_t = 2.11$ min

5 MS (ESIpos): $m/z = 611$ ($M+H$)⁺

¹H-NMR (300 MHz, CDCl₃): $\delta = 7.75$ - 7.20 (m, 8H), 6.28 (s, 1H), 5.96 (d, 1H), 5.62 (d, 1H), 4.72 (d, 1H), 4.55 (d, 1H), 1.47-1.35 (m, 15H) ppm.

Example 30

10 [6-(4-Cyanophenyl)-5-(cyclobutylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2H)-yl]acetic acid



15 *tert.*-Butyl [6-(4-cyanophenyl)-5-(cyclobutylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidine-1(2H)-yl]acetate (Example 26) (35 mg, 0.063 mmol) is dissolved in trifluoroacetic acid (1 ml). After 5 minutes stirring, the solution is concentrated *in vacuo* and the residue is purified by HPLC (RP18 column; eluent: acetonitrile/0.1% aq. formic acid 10:90 → 90:10).

Yield: 23.7 mg (65% of th.)

LC-MS (method 4): $R_t = 2.51$ min


HPLC (method 1): $R_t = 4.82$ min, $\lambda_{\max} = 234$ nm

MS (ESIpos): $m/z = 498 (M+H)^+$

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6): $\delta = 12.5$ (br s, 1H), 7.96-7.54 (m, 8H), 5.61 (s, 1H), 4.19 (d, 1H), 3.85 (d, 1H), 2.30-1.60 (m, 7H), 1.91 (s, 3H) ppm.

Example 31

- 5 [6-(4-Cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetic acid



The title compound is prepared from Example 27 according to the procedure described for Example 30.

- 10 Yield: 27.1 mg (35% of th.)

LC-MS (method 3): $R_t = 2.20$ min


HPLC (method 6): $R_t = 4.47$ min, $\lambda_{\text{max}} = 234$ nm

MS (ESIpos): $m/z = 484 (M+H)^+$

- 15 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6): $\delta = 13.0$ (br s, 1H), 7.93-7.53 (m, 8H), 5.76 (s, 1H), 4.23-3.71 (m, 2H), 2.26 (m, 1H), 1.98 (s, 3H), 0.96-0.73 (m, 4H) ppm.

Example 32

[6-(4-Cyanophenyl)-5-(cyclohexylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidine-1(2*H*)-yl]acetic acid



The title compound is prepared from Example 28 according to the procedure described for Example 30.

Yield: 18 mg (39% of th.)

5 LC-MS (method 4): $R_t = 2.69$ min


HPLC (method 6): $R_t = 5.01$ min, $\lambda_{\max} = 236$ nm

MS (ESIpos): $m/z = 526$ ($M+H$)⁺

¹H-NMR (300 MHz, DMSO-*d*₆): $\delta = 12.5$ (br s, 1H), 7.99-7.50 (m, 8H), 5.70 (s, 1H), 4.25-3.74 (m, 2H), 2.75 (m, 1H), 1.85 (s, 3H), 1.78-0.80 (m, 10H) ppm.

10 *Example 33*

tert.-Butyl [6-(4-cyanophenyl)-5-(4-methoxybenzoyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)-phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate



The title compound is prepared from Example 24 according to the procedure described for Example 26.

Yield: 141.2 mg (76% of th.)


5 HPLC (method 6): $R_t = 5.36$ min, $\lambda_{\max} = 189$ nm

MS (ESIpos): $m/z = 606$ (M+H)⁺

¹H-NMR (300 MHz, CDCl₃): $\delta = 7.79$ - 7.40 (m, 8H), 7.38 (d, 2H), 6.92 (d, 2H), 5.62 (s, 1H), 4.61 (d, 1H), 3.88 (s, 3H), 3.39 (d, 1H), 1.55 (s, 3H), 1.49 (s, 9H) ppm.

Example 34

10 [6-(4-Cyanophenyl)-5-(4-methoxybenzoyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidine-1(2H)-yl]acetic acid



The title compound is prepared from Example 33 according to the procedure described for Example 30.

Yield: 81 mg (81% of th.)

LC-MS (method 3): $R_t = 2.34$ min


5 HPLC (method 6): $R_t = 4.75$ min, $\lambda_{\max} = 200$ nm

MS (ESIpos): $m/z = 550$ (M+H)⁺

¹H-NMR (300 MHz, DMSO-d₆): $\delta = 12.72$ (br s, 1H), 7.96-7.54 (m, 10H), 6.98 (d, 2H), 5.66 (s, 1H), 4.21 (d, 1H), 3.81 (s, 3H), 3.19 (d, 1H), 1.41 (s, 3H) ppm.

Example 35

10 *tert.*-Butyl 3-[[6-(4-cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoro-methyl)-phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]methyl}benzoate



15 Sodium hydride (12.1 mg, 0.31 mmol; 60% suspension in mineral oil) is washed with pentane (3 x 3 ml), then treated with a solution of 4-{5-(cyclopropylcarbonyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl}benzonitrile (Example 22) (100 mg, 0.235 mmol) in tetrahydrofuran (5 ml). After 5 minutes, the reaction is treated with a solution of *tert.*-butyl 3-(bromomethyl)benzoate (77 mg, 0.28 mmol) in tetrahydrofuran (5 ml) and then stirred at room temperature overnight (16 h). The reaction mixture is quenched with water (50 ml) and extracted with ethyl acetate (3 x 150 ml). The combined organic phases are washed with brine, dried over

anhydrous magnesium sulfate, filtered and concentrated *in vacuo*. The crude product is purified by preparative HPLC (RP18 column; eluent: acetonitrile/0.1% aq. formic acid 10:90 → 90:10).

Yield: 124 mg (84% of th.)

LC-MS (method 3): $R_t = 3.04$ min


5 HPLC (method 6): $R_t = 5.59$ min, $\lambda_{\max} = 198$ nm

MS (ESIpos): $m/z = 616$ (M+H)⁺

¹H-NMR (300 MHz, CDCl₃): $\delta = 7.93$ -7.83 (m, 2H), 7.73-7.58 (m, 4H), 7.54-7.35 (m, 6H), 5.50 (s, 1H), 5.21 (d, 1H), 4.00 (d, 1H), 2.01 (s, 3H), 1.94 (m, 1H), 1.60 (s, 9H), 1.08-0.79 (m, 4H) ppm.

Example 36

10 3-{[6-(4-Cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2H)-yl]methyl} benzoic acid



The title compound is prepared from Example 35 according to the procedure described for Example 30, with the exception that the reaction time is 30 minutes, and the title compound is
15 purified by preparative HPLC (RP18 column; eluent: acetonitrile/water 30:70 → 90:10).

Yield: 67 mg (82% of th.)


LC-MS (method 4): $R_t = 2.59$ min

MS (ESIpos): $m/z = 560$ (M+H)⁺

¹H-NMR (300 MHz, DMSO-d₆): δ = 12.90 (br s, 1H), 7.86-7.35 (m, 12H), 5.68 (s, 1H), 4.90 (d, 1H), 4.30 (d, 1H), 2.21 (m, 1H), 1.95 (s, 3H), 0.95-0.76 (m, 4H) ppm.

Example 37

tert.-Butyl 4-({4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetra-
5 hydropyrimidin-5-yl}carbonyl)piperidine-1-carboxylate



The title compound is prepared according to the procedure described for Example 18.

Yield: 20 mg (2% of th.)

LC-MS (method 4): R_t = 2.73 min


10 HPLC (method 6): R_t = 5.09 min, λ_{max} = 196 nm

MS (ESIpos): m/z = 569 (M+H)⁺

¹H-NMR (300 MHz, DMSO-d₆): δ = 8.40 (br d, 1H), 8.17-7.19 (m, 8H), 5.51 (br d, 1H), 4.00-3.70 (m, 1H), 3.03-2.39 (m, 3H), 1.85 (s, 3H), 1.80-1.66 (m, 2H), 1.37 (s, 9H), 1.56-1.00 (m, 2H) ppm.

Example 38

15 *tert.*-Butyl 3-{{[6-(4-cyanophenyl)-5-(cyclohexylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2H)-yl]methyl}benzoate



The title compound is prepared from Example 21 according to the procedure described for Example 35.

Yield: 65 mg (46% of th.)


5 LC-MS (method 3): $R_t = 3.31$ min

MS (ESIpos): $m/z = 658$ ($M+H$)⁺

¹H-NMR (200 MHz, DMSO- d_6): $\delta = 7.90$ - 7.39 (m, 12 H), 5.58 (s, 1H), 4.97 (d, 1H), 4.31 (d, 1H), 2.67 (m, 1H), 1.80 (s, 3H), 1.52 (s, 9H), 1.70-0.85 (m, 10 H) ppm.

Example 39

10 *tert.*-Butyl 4-({3-(2-*tert.*-butoxy-2-oxoethyl)-4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl} carbonyl)piperidine-1-carboxylate



A stirred suspension of *tert.*-butyl 4-({4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl}carbonyl)piperidine-1-carboxylate (Example 37) (43 mg, 0.08 mmol) and potassium carbonate (19 mg, 0.14 mmol) in dimethylformamide (1.5 ml) is treated with *tert.*-butyl bromoacetate (16 mg, 0.08 mmol), then stirred at room temperature overnight (16 h). The reaction solution is then diluted with methanol (7 ml) and purified directly by preparative HPLC (RP18 column; eluent: acetonitrile/water 10:90 → 90:10).

Yield: 26.5 mg (51% of th.)

LC-MS (method 3): $R_t = 3.02$ min


MS (ESIpos): $m/z = 682$ (M+H)⁺

10 HPLC (method 6): $R_t = 5.44$ min, $\lambda_{\max} = 234$ nm

¹H-NMR (300 MHz, DMSO-*d*₆): $\delta = 8.43$ -7.34 (m, 8H), 5.71 (s, 1H), 4.22-3.74 (m, 4H), 3.10-2.39 (m, 4H), 1.84 (s, 3H), 1.88-1.69 (m, 1H), 1.54-1.00 (s, 20H) ppm.

Example 40

3-{{[6-(4-Cyanophenyl)-5-(cyclohexylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]methyl}benzoic acid



The title compound is prepared from Example 38 according to the procedure described for Example 30, with the exception that the reaction time is 30 minutes, and the title compound is purified by preparative HPLC (RP18 column; eluent: acetonitrile/water 30:70 → 90:10).

20 Yield: 40 mg (77% of th.)


HPLC (method 6): $R_t = 5.15$ min, $\lambda_{\max} = 200$ nm

MS (ESIpos): $m/z = 602$ (M+H)⁺

¹H-NMR (300 MHz, DMSO-d₆): $\delta = 12.95$ (s, 1H), 7.94-7.40 (m, 12H), 5.51 (s, 1H), 5.00 (d, 1H), 4.21 (d, 1H), 2.77-2.60 (m, 1H), 1.80 (s, 3H), 1.68-1.36 (m, 5H), 1.29-0.90 (m, 5H) ppm.

5 Example 41

Diethyl {4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl}phosphonate



10 A solution of diethyl (2-oxopropyl)phosphonate (250 mg, 1.29 mmol), 4-cyanobenzaldehyde (168.84 mg, 1.29 mmol), *N*-[3-(trifluoromethyl)phenyl]urea (263 mg, 1.29 mmol) and polyphosphoric acid ethyl ester (0.70 g) in tetrahydrofuran (5 ml) is refluxed overnight with stirring. The reaction solution is purified directly by preparative HPLC (RP18 column; eluent: acetonitrile/water 10:90 → 90:10). The fractions containing product are concentrated *in vacuo* and re-purified by flash chromatography on silica gel 60 with dichloromethane / methanol as eluent.

15 Yield: 20 mg (3% of th.)


HPLC (method 1): $R_t = 4.31$ min, $\lambda_{\max} = 236$ nm

MS (ESIpos): $m/z = 494$ (M+H)⁺

¹H-NMR (300 MHz, DMSO-d₆): $\delta = 8.39$ -8.27 (m, 1H), 7.07-7.48 (m, 8H), 4.98-4.84 (m, 1H), 3.79-3.19 (m, 4H), 2.50 (s, 3H), 1.05 (t, 3H), 0.95 (t, 3H) ppm.

Example 42

Methyl 5-{{[6-(4-cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)-phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]methyl}-2-furoate



- 5 To a stirred suspension of 4-{5-(cyclopropylcarbonyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)-phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl} benzonitrile (Example 22) (150 mg, 0.35 mmol) and potassium carbonate (98 mg, 0.71 mmol) in dimethylformamide (3 ml) is added methyl 5-(chloromethyl)-2-furoate (92 mg, 0.53 mmol). The suspension is stirred at room temperature overnight (16 h), then additional methyl 5-(chloromethyl)-2-furoate (6.1 mg, 0.35 mmol) and potassium carbonate (49 mg, 0.35 mmol) are added, and the suspension is stirred an additional 72 hours. The reaction mixture is diluted with methanol (5 ml) and purified directly by preparative HPLC (RP18 column; eluent: acetonitrile/water 10:90 → 90:10). The title compound is isolated as a brownish amorphous solid.

Yield: 72 mg (35% of th.)

- 15 LC-MS (method 4): $R_t = 2.75$ min


MS (ESIpos): $m/z = 564$ ($M+H$)⁺

HPLC (method 1): $R_t = 4.98$ min, $\lambda_{max} = 196$ nm

- ¹H-NMR (300 MHz, DMSO-*d*₆): $\delta = 7.99$ -7.41 (m, 8H), 7.13 (d, 1H, $J = 3.58$ Hz), 6.47 (d, 1H, $J = 3.58$ Hz), 5.73 (s, 1H), 4.78 (d, 1H), 4.50 (d, 1H), 3.78 (s, 3H), 2.57-2.48 (m, 1H), 1.94 (s, 3H), 1.03-0.73 (m, 4H) ppm.
- 20

Example 43

Methyl 2-{[6-(4-cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)-phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]methyl}-1,3-oxazole-4-carboxylate



5 To a stirred suspension of 4-{5-(cyclopropylcarbonyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)-phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl} benzonitrile (Example 22) (150 mg, 0.35 mmol) and potassium carbonate (98 mg, 0.71 mmol) in dimethylformamide (3 ml) is added methyl 2-(chloromethyl)-1,3-oxazole-4-carboxylate (93 mg, 0.53 mmol). The reaction mixture is stirred at room temperature overnight, then an additional equivalent of potassium carbonate (0.35 mmol) and

10 methyl 2-(chloromethyl)-1,3-oxazole-4-carboxylate (0.35 mmol) are added, and the solution is stirred an additional 72 hours. The reaction mixture is quenched with water (20 ml) and extracted with ethyl acetate (3 x 50 ml). The combined organic phases are washed with brine, dried over anhydrous magnesium sulfate, filtered and concentrated. The residue is purified by HPLC (RP18 column; eluent: acetonitrile/water 30:70 → 90:10).

15 Yield: 74 mg (37% of th.)

LC-MS (method 5): $R_t = 2.61$ min

MS (ESIpos): $m/z = 565$ ($M+H$)⁺


HPLC (method 1): $R_t = 4.78$ min, $\lambda_{max} = 196$ nm

¹H-NMR (300 MHz, DMSO-*d*₆): $\delta = 8.68$ (s, 1H), 7.92-7.48 (m, 8H), 5.84 (s, 1H), 4.91-4.51 (m, 2H), 3.79 (s, 3H), 2.37-2.23 (m, 1H), 1.96 (s, 3H), 0.96-0.72 (m, 4H) ppm.

20

Example 44

2-[6-(4-Cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]-*N*-[(4-cyanophenyl)sulfonyl]acetamide



- 5 A mixture of [6-(4-cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)-phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetic acid (Example 31) (100 mg, 0.21 mmol), 1,3-dicyclohexylcarbodiimide (41.5 mg, 0.23 mmol), 4-cyanobenzene-1-sulfonamide (41.5 mg, 0.23 mmol) and 4-dimethylaminopyridine (28 mg, 0.23 mmol) in dichloromethane (4 ml) is stirred for 48 hours. The product is extracted with dichloromethane, washed with 2 N hydrochloric acid and
- 10 brine, dried over anhydrous magnesium sulfate, filtered and concentrated. The residue is purified by preparative HPLC (RP18 column; eluent: acetonitrile/water 10:90 → 90:10).

Yield: 45 mg (33% of th.)

LC-MS (method 3): $R_t = 2.56$ min


MS (ESIpos): $m/z = 648$ ($M+H$)⁺

- 15 HPLC (method 1): $R_t = 4.83$ min, $\lambda_{max} = 198$ nm

¹H-NMR (300 MHz, DMSO-*d*₆): $\delta = 8.22$ -7.98 (m, 4H), 7.93-7.45 (m, 9H), 5.65 (s, 1H), 4.30-3.63 (m, 2H), 2.34-2.13 (m, 1H), 1.94 (s, 3H), 0.98-0.70 (m, 4H) ppm.

Example 45

- 2-[6-(4-Cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-
- 20 dihydropyrimidin-1(2*H*)-yl]-*N*-[(2,2,2-trifluoroethyl)sulfonyl]acetamide



A mixture of [6-(4-cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)-phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetic acid (Example 31) (100 mg, 0.21 mmol), 2,2,2-trifluoroethanesulfonamide (37 mg, 0.23 mmol), 1,3-dicyclohexylcarbodiimide (47 mg, 0.23 mmol) and 4-dimethylaminopyridine (28 mg, 0.23 mmol) in dichloromethane (4 ml) is stirred for 48 hours. The crude product is extracted with dichloromethane (100 ml), washed with 2 N hydrochloric acid and brine, dried over anhydrous magnesium sulfate, filtered and concentrated. The residue is purified by flash chromatography with silica gel 60 (50 g) and 2:1 cyclohexane/ethyl acetate as eluent.

Yield: 73 mg (56% of th.)

MS (ESIpos): $m/z = 629$ ($M+H$)⁺


HPLC (method 1): $R_t = 4.84$ min, $\lambda_{max} = 234$ nm

¹H-NMR (300 MHz, DMSO-*d*₆): $\delta = 7.94$ - 7.48 (m, 9H), 5.75 (s, 1H), 4.3 (d, 1H), 4.08 (q, 2H), 3.12 (d, 1H), 2.44-2.25 (m, 1H), 1.90 (s, 3H), 0.99-0.72 (m, 4H) ppm.

Example 46

Methyl 5-{{[6-(4-cyanophenyl)-5-(cyclohexylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)-phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]methyl}-2-furoate

81



The title compound is prepared from Example 21 according to the procedure described for Example 42, with the exception that only 1.5 equivalents of methyl 5-(chloromethyl)-2-furoate and 2 equivalents of potassium carbonate are used. The title compound is purified by preparative
 5 HPLC (RP18 column; eluent: acetonitrile/0.1% aq. formic acid 30:70 → 90:10).

Yield: 65 mg (34% of th.)


LC-MS (method 4): $R_t = 3.10$ min

MS (ESIpos): $m/z = 606$ ($M+H$)⁺

¹H-NMR (300 MHz, DMSO- d_6): $\delta = 7.95$ - 7.48 (m, 8H), 7.19 (m, 1H), 6.53 (m, 1H), 5.59 (s, 1H),
 10 4.88 (d, 1H), 4.43 (d, 1H), 3.79 (s, 3H), 2.78-2.65 (m, 1H), 1.78 (s, 3H), 1.70-0.97 (m, 10H).

Example 47

Methyl 2-{[6-(4-cyanophenyl)-5-(cyclohexylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)-phenyl]-3,6-dihydropyrimidin-1(2H)-yl]methyl}-1,3-oxazole-4-carboxylate



The title compound is prepared from Example 21 according to the procedure described for Example 43, with the exception that only 2 equivalents of potassium carbonate and 1.5 equivalents of methyl 2-(chloromethyl)-1,3-oxazole-4-carboxylate are used. The reaction time is 72 hours.

5 Yield: 80 mg (41% of th.)

LC-MS (method 4): $R_t = 2.76$ min


MS (ESIpos): $m/z = 607$ ($M+H$)⁺

HPLC (method 1): $R_t = 5.41$ min, $\lambda_{\max} = 194$ nm

¹H-NMR (300 MHz, DMSO-*d*₆): $\delta = 8.71$ (s, 1H), 7.93-7.54 (m, 8H), 5.74 (s, 1H), 4.93 (d, 1H),
 10 4.53 (d, 1H), 3.80 (s, 3H), 2.79-2.67 (m, 1H), 1.81 (s, 3H), 1.69-0.97 (m, 10H) ppm.

Example 48

5-{[6-(4-Cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]methyl}-2-furoic acid



A stirred solution of methyl 5-[[6-(4-cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]methyl]-2-furoate (Example 42) (50 mg, 0.09 mmol) in tetrahydrofuran (1.5 ml) is treated with a solution of lithium hydroxide (2.34 mg, 0.1 mmol) in water (1.5 ml). After three hours stirring at room temperature, additional lithium hydroxide (2.34 mg, 0.1 mmol) is added. The reaction solution is stirred overnight (16 h), then allowed to stand for 48 h. The solution is acidified with 1 N hydrochloric acid (500 μ l). After 5 minutes stirring, a precipitate is obtained. Methanol (3 ml) is added, and the crude reaction solution is purified directly by preparative HPLC (RP18 column; eluent: acetonitrile/water 30:70 \rightarrow 90:10). The title compound is isolated as a brownish solid.

Yield: 27.5 mg (56.4% of th.)


HPLC (method 6): $R_t = 4.71$ min, $\lambda_{\max} = 244$ nm

MS (ESIpos): $m/z = 550$ ($M+H$)⁺

¹H-NMR (400 MHz, DMSO- d_6): $\delta = 13.1$ (br s, 1H), 7.90-7.76 (m, 4H), 7.74 (t, 1H), 7.63 (d, 1H), 7.51 (d, 2H), 7.03 (m, 1H), 6.45 (d, 1H), 5.72 (s, 1H), 4.84 (d, 1H), 4.38 (d, 1H), 2.28 (m, 1H), 1.95 (s, 3H), 0.98-0.75 (m, 4H) ppm.

Example 49

2-[6-(4-Cyanophenyl)-5-(cyclohexylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]-*N*-[(4-cyanophenyl)sulfonyl]acetamide



The title compound is prepared from Example 32 according to the procedure described for Example 44, with the exception that the title compound is purified by HPLC (RP18 column; eluent: acetonitrile/water 30:70 → 90:10).

5 Yield: 65 mg (48% of th.)

LC-MS (method 3): $R_t = 2.93$ min


MS (ESIpos): $m/z = 690$ ($M+H$)⁺

HPLC (method 1): $R_t = 5.28$ min, $\lambda_{max} = 198$ nm

10 ¹H-NMR (300 MHz, DMSO-*d*₆): $\delta = 12.57$ (br s, 1H), 8.06 (d of d, 4H), 7.84 (d, 2H), 7.75-7.62 (m, 3H), 7.58-7.49 (m, 3H), 5.56 (s, 1H), 4.16 (d, 1H), 3.86 (d, 1H), 2.64 (m, 1H), 1.79 (s, 3H), 1.68-0.93 (m, 10H) ppm.

Example 50

2-[6-(4-Cyanophenyl)-5-(cyclohexylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2H)-yl]-*N*-[(2,2,2-trifluoroethyl)sulfonyl]acetamide



The title compound is prepared from Example 32 according to the procedure described for Example 45, with the exception that the title compound is purified by preparative HPLC (RP18 column; eluent: acetonitrile/water 30:70 → 90:10).

5 Yield: 52 mg (40% of th.)


HPLC (method 1): $R_t = 5.30$ min, $\lambda_{\max} = 238$ nm

MS (ESIpos): $m/z = 671$ ($M+H$)⁺

¹H-NMR (300 MHz, DMSO-*d*₆): $\delta = 7.89$ (d, 2H), 7.80 (d, 1H), 7.71 (t, 2H), 7.62 (d, 2H), 7.59 (s, 1H), 5.66 (s, 1H), 4.70-4.50 (m, 2H), 4.28 (d, 1H), 3.86 (d, 1H), 2.72 (m, 1H), 1.82 (s, 3H), 1.91-
10 0.96 (m, 10H) ppm.

Example 51

2-{{[6-(4-Cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]methyl}-1,3-oxazole-4-carboxylic acid



To a stirred solution of methyl 2-{{[6-(4-cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2H)-yl]methyl}-1,3-oxazole-4-carboxylate (Example 43) (50 mg, 0.09 mmol) in tetrahydrofuran (1.5 ml) is added a solution of lithium hydroxide (4.24 mg, 0.18 mmol) in water (1.5 ml). The reaction solution is stirred at room temperature overnight (16 h). The pH of the solution is adjusted to less than pH 7 with 1 N hydrochloric acid (500 µl). After 5 minutes stirring, a precipitate is obtained. Methanol (3 ml) is added, and the crude reaction solution is purified directly by preparative HPLC (RP18 column; eluent: acetonitrile/0.1% aq. formic acid 30:70 → 90:10). The title compound is isolated as a colourless solid.

Yield: 41 mg (84% of th.)


LC-MS (method 3): $R_t = 2.22$ min

MS (ESIpos): $m/z = 551$ ($M+H$)⁺

¹H-NMR (300 MHz, DMSO- d_6): $\delta = 13.01$ (br s, 1H), 8.57 (s, 1H), 7.90-7.52 (m, 8H), 5.84 (s, 1H), 4.88 (d, 1H), 4.51 (d, 1H), 2.30 (m, 1H), 1.96 (s, 3H), 0.96-0.73 (m, 4H) ppm.

Example 52

5-{{[6-(4-Cyanophenyl)-5-(cyclohexylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2H)-yl]methyl}-2-furoic acid



The title compound is prepared from Example 46 according to the procedure described for Example 48, with the exception that the title compound is purified by preparative HPLC (RP18 column; eluent: acetonitrile/0.1% aq. formic acid 10:90 → 90:10).

5 Yield: 43 mg (77% of th.)

HPLC (method 1): $R_t = 5.20$ min, $\lambda_{\max} = 202$ nm


LC-MS (method 4): $R_t = 2.80$ min

MS (ESIpos): $m/z = 592$ ($M+H$)⁺

10 ¹H-NMR (300 MHz, DMSO- d_6): $\delta = 13.0$ (br s, 1H), 7.90-7.65 (m, 5H), 7.63-7.46 (m, 3H), 7.10 (d, 1H), 6.53 (d, 1H), 5.55 (s, 1H), 4.92 (d, 1H), 4.32 (d, 1H), 2.77-2.63 (m, 1H), 1.79 (s, 3H), 1.68-1.37 (m, 5H), 1.31-0.93 (m, 5H) ppm.

Example 53

2-([6-(4-Cyanophenyl)-5-(cyclohexylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2H)-yl]methyl)-1,3-oxazole-4-carboxylic acid



The title compound is prepared from Example 47 according to the procedure described for Example 51, with the exception that the title compound is purified by preparative HPLC (RP18 column; eluent: acetonitrile/0.1% aq. formic acid 10:90 → 90:10).

5 Yield: 46 mg (84% of th.)

LC-MS (method 3): $R_t = 2.52$ min


MS (ESIpos): $m/z = 593$ ($M+H$)⁺

HPLC (method 1): $R_t = 5.07$ min, $\lambda_{\max} = 194$ nm

10 ¹H-NMR (300 MHz, DMSO-d₆): $\delta = 13.0$ (br s, 1H), 8.58 (s, 1H), 7.86 (d, 2H), 7.83-7.60 (m, 4H), 7.58 (d, 2H), 5.72 (s, 1H), 4.95 (d, 1H), 4.47 (d, 1H), 2.80-2.66 (m, 1H), 1.81 (s, 3H), 1.69-0.95 (m, 10H) ppm.

Example 54

Benzyl 2-(4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl}carbonyl)piperidine-1-carboxylate



A stirred mixture of benzyl 2-acetoacetyl piperidine-1-carboxylate (Example 10A) (1.74 g, 5.74 mmol), 4-cyanobenzaldehyde (752 mg, 5.74 mmol), *N*-[3-(trifluoromethyl)phenyl]urea (976 mg, 4.78 mmol) and 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide anhydride (6.1 g, 9.5 mmol) in methyl *tert*.-butyl ether (44 ml) is refluxed overnight (16 h). The product is extracted with methyl *tert*.-butyl ether (300 ml), washed with aq. saturated sodium bicarbonate solution (200 ml), dried with anhydrous magnesium sulfate, filtered and concentrated. The residue is chromatographed with silica gel 60 using cyclohexane / ethyl acetate as eluent.

Yield: 858 mg (30% of th.)

10 HPLC (method 1): $R_t = 5.16$ min, $\lambda_{\max} = 232$ nm


MS (ESIpos): $m/z = 603$ ($M+H$)⁺

LC-MS (method 3): $R_t = 2.66$ min

¹H-NMR (300 MHz, CDCl₃): $\delta = 7.8-7.1$ (m, 14H), 5.80-4.80 (m, 3H), 4.2-3.8 (m, 1H), 3.3-2.9 (m, 2H), 1.5 (s, 3H), 2.1-0.8 (m, 6H) ppm.

15 Example 55

Benzyl 2-({3-(2-*tert*.-butoxy-2-oxoethyl)-4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl}carbonyl)piperidine-1-carboxylate



The title compound is prepared from Example 54 according to the procedure described for Example 26.

Yield: 53 mg (18% of th.)

5 LC-MS (method 3): $R_t = 3.08$ min


HPLC (method 1): $R_t = 5.62$ min, $\lambda_{\max} = 234$ nm

MS (ESIpos): $m/z = 712$ (M+H)⁺

¹H-NMR (300 MHz, CDCl₃): $\delta = 7.72$ - 7.12 (m, 13H), 5.64 - 4.56 (m, 5H), 4.18 - 3.92 (m, 1H), 3.52 - 3.12 (m, 2H), 2.35 - 0.73 (m, 6H), 1.52 (s, 3H), 1.48 (s, 9H) ppm.

10 Example 56

[5-({1-[(Benzyloxy)carbonyl]piperidin-2-yl}carbonyl)-6-(4-cyanophenyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2H)-yl]acetic acid



The title compound is prepared from Example 55 according to the procedure described for Example 30.


Yield: 35 mg (84% of th.)

- 5 HPLC (method 1): $R_t = 5.70$ min, $\lambda_{\max} = 254$ nm

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6): $\delta = 12.5$ (br s, 1H), 7.91-7.01 (m, 13H), 5.86 (d, 1H), 5.40-5.25 (m, 1H), 5.08 (s, 1H), 5.00-4.76 (m, 1H), 4.27 (m, 1H), 3.93-3.68 (m, 1H), 1.75 (s, 3H), 1.9-0.7 (m, 8H) ppm.


In analogy to the procedure for Example 2A, the following compounds are prepared:

Example No.	Structure	Starting material	Yield [%]	R_t [min] (method)	Mass $[\text{M}+\text{H}]^+$
57		Example 11A	72	5.50 (1)	656

Example No.	Structure	Starting material	Yield [%]	R _t [min] (method)	Mass [M+H] ⁺
58		Example 12A	13	5.22 (6)	564

Example 59

Benzyl *tert*.-butyl ({4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl}carbonyl)malonate



5

1401 mg (5.60 mmol) benzyl *tert*.-butyl malonate are dissolved in 4 ml tetrahydrofuran under an argon atmosphere. The solution is cooled to 0°C, 156 mg (3.92 mmol) sodium hydride are added, and the mixture is warmed to room temperature and stirred for 30 minutes. 580 mg (1.12 mmol) of Example 16A are added as a solution in 6 ml tetrahydrofuran. The reaction mixture is stirred under reflux overnight. The mixture is partitioned between ethyl acetate and 2 N hydrochloric acid, the organic phase is sequentially washed with water and saturated sodium chloride solution, dried over magnesium sulfate and evaporated *in vacuo*. The crude product is purified by column chromatography.

10


graphy on silica gel (eluent: cyclohexane/ethyl acetate 2:1 → 0:1) and thereafter by RP-HPLC with a water/acetonitrile gradient.

Yield: 270 mg (38% of th.) as a mixture of diastereomers

¹H-NMR (300 MHz, DMSO-d₆): δ = 1.21/1.36 (2s, 9H), 1.88/1.92 (2s, 3H), 4.95 (s, 1H), 5.10-5.23 (m, 2H), 5.47/5.51 (2d, 1H), 7.21-7.90 (m, 13H), 8.58/8.61 (2d, 1H) ppm.

Example 60

Benzyl 3-{4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl}-3-oxopropanoate




10 100 mg (0.16 mmol) of Example 59 are dissolved in 2 ml dichloromethane/trifluoroacetic acid (1:1) and stirred at room temperature for 1 hour. Volatiles are evaporated *in vacuo* and the crude product is purified by column chromatography on silica gel (eluent: cyclohexane/ethyl acetate 1:1).

Yield: 39 mg (42% of th.)

¹H-NMR (400 MHz, DMSO-d₆): δ = 1.99 (s, 3H), 3.58 (d, 1H), 3.93 (d, 1H), 5.05 (s, 2H), 5.48 (d, 1H), 7.25-7.40 (m, 4H), 7.48-7.55 (br d, 1H), 7.60-7.73 (m, 4H), 7.80 (d, 1H), 7.86 (d, 2H), 8.55 (d, 1H) ppm.

Example 61

2-(Benzyloxy)-2-oxoethyl (4*R*)-4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidine-5-carboxylate



200 mg (0.50 mmol) of Example 6A are dissolved in 2.0 ml dimethylformamide. 456 mg (1.99 mmol) benzyl bromoacetate and 100 mg (1.00 mmol) triethylamine are added, and the mixture is stirred at room temperature for 2 hours. The reaction mixture is partitioned between
 5 ethyl acetate and 2 N hydrochloric acid, the organic layer is sequentially washed with water and saturated sodium chloride solution, dried over magnesium sulfate and evaporated to dryness *in vacuo*. The crude product is purified by column chromatography on silica gel (eluent: cyclohexane/ethyl acetate 1:1 → 0:1).

Yield: 260 mg (95% of th.)

10 ¹H-NMR (200 MHz, DMSO-d₆): δ = 2.05 (s, 3H), 4.74 (s, 1H), 5.12 (s, 1H), 5.41 (d, 1H), 7.28-7.42 (m, 5H), 7.53 (d, 1H), 7.61 (d, 2H), 7.65-7.75 (m, 2H), 7.77-7.86 (m, 3H), 8.45 (d, 1H) ppm.


In analogy to the procedure for Example 13A, the following compounds are prepared:

Example No.	Structure	Starting material	Yield [%]	R _t [min] (method)	Mass [M+H] ⁺
62		Example 57	60	5.91 (1)	770

Example No.	Structure	Starting material	Yield [%]	R _t [min] (method)	Mass [M+H] ⁺
63		Example 58	62	5.58 (1)	678

Example 64

2-[(*tert.*-Butoxycarbonyl)amino]ethyl 3-(2-*tert.*-Butoxy-2-oxoethyl)-4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidine-5-carboxylate



5


70 mg (0.12 mmol) of Example 15A and 521 mg (3.23 mmol) *tert.*-butyl-*N*-hydroxyethyl carbamate are reacted at 100°C for 150 minutes. The reaction mixture is diluted with acetonitrile and purified using preparative RP-HPLC (eluent: acetonitrile/water gradient).

Yield: 41.9 mg (52% of th.)

10 ¹H-NMR (200 MHz, DMSO-*d*₆): δ = 1.28 (s, 9H), 1.36 (s, 9H), 2.05 (s, 3H), 3.03-3.18 (m, 2H), 3.80-4.1 (m, 4H), 5.10 (s, 1H), 6.82-6.95 (m, 1H), 7.60-7.93 (m, 8H) ppm.

Example 65

2-(2-Oxopyrrolidin-1-yl)ethyl 3-(2-*tert.*-butoxy-2-oxoethyl)-4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidine-5-carboxylate




5 The title compound is synthesized in analogy to Example 64 from Example 15A using *N*-(2-hydroxyethyl)-2-pyrrolidone instead of *tert*.-butyl-*N*-hydroxyethyl carbamate.

Yield: 74% of th.

¹H-NMR (200 MHz, DMSO-d₆): δ = 1.30 (s, 9H), 1.81 (q, 2H), 2.03 (s, 3H), 2.13 (t, 2H), 3.13-3.5 (m, 4H), 3.84 (d, 1H), 4.00-4.20 (m, 3H), 5.52 (s, 1H), 7.60-7.92 (m, 8H) ppm.

10 *Example 66*

2-(Benzyloxy)-2-oxoethyl 3-(2-*tert.*-butoxy-2-oxoethyl)-4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidine-5-carboxylate




The title compound is synthesized in analogy to Example 64 from Example 15A using benzyl hydroxyacetate instead of *tert*.-butyl-*N*-hydroxyethyl carbamate.

Yield: 32% of th.

- 5 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6): δ = 1.30 (s, 9H), 2.05 (s, 3H), 3.87 (d, 1H), 4.09 (d, 1H), 4.73 (d, 1H), 5.12 (s, 1H), 5.60 (s, 1H), 7.28-7.41 (m, 5 H), 7.60-7.85 (m, 8H) ppm.

Example 67

[6-(4-Cyanophenyl)-4-methyl-2-oxo-5-{[2-(2-oxopyrrolidin-1-yl)ethoxy]carbonyl}-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetic acid




34.9 mg of Example 65 are dissolved in 2 ml dichloromethane/trifluoroacetic acid (1:1) and stirred at room temperature overnight. Volatiles are evaporated *in vacuo* and the remainder is purified by RP-HPLC using a water/acetonitrile gradient.

Yield: 18 mg (57% of th.)

- 5 $^1\text{H-NMR}$ (200 MHz, DMSO-d_6): δ = 1.81 (q, 2H), 2.02 (s, 3H), 2.13 (t, 2H), 3.10-3.55 (m, 4H), 3.72 (d, 1H), 4.00-4.22 (m, 3H), 5.52 (s, 1H), 7.58-7.93 (m, 8H), 12.75 (br s, 1H) ppm.

Example 68

tert.-Butyl [6-(4-cyanophenyl)-4-methyl-5-[(methylsulfonyl)amino]carbonyl]-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate



10



- 100 mg (0.19 mmol) of Example 14A, 36.9 mg (0.39 mmol) methanesulfonamide, 44.0 mg (0.21 mmol) dicyclohexylcarbodiimide and 26.1 mg (0.21 mmol) 4-*N,N*-dimethylaminopyridine are dissolved in dry dichloromethane (2 ml) and reacted overnight. The reaction mixture is filtered, the filtrate is sequentially washed with 2 M hydrochloric acid, water and saturated ammonium chloride solution, and the organic phase is dried over magnesium sulfate and evaporated to dryness. The crude product is purified by column chromatography on silica gel (eluent: cyclohexane/ethyl acetate/acetic acid 50:50:1).
- 15

Yield: 50 mg (41% of th.)

- $^1\text{H-NMR}$ (200 MHz, DMSO-d_6): δ = 1.28 (s, 3H), 1.80 (s, 3H), 3.07 (s, 3H), 3.80 (d, 1H), 4.01 (d, 1H), 5.60 (s, 1H), 7.56-7.95 (m, 8H), 11.70 (br s, 1H) ppm.
- 20


In analogy to the procedure for Example 68, the following compounds are prepared:

Example No.	Structure	Starting materials	Yield [%]	R _t [min] (method)	Mass [M+H] ⁺
69		Example 14A; 2,2,2-trifluoroethanesulfonamide	71	5.12 (1)	661
70		Example 14A; 4-fluorobenzene-sulfonamide	43	5.17 (1)	673
71		Example 5A; methanesulfonamide	6	4.17 (6)	479

Example No.	Structure	Starting materials	Yield [%]	R _t [min] (method)	Mass [M+H] ⁺
72		Example 5A; 2,2,2-trifluoroethanesulfonamide	76	4.51 (6)	547
73		Example 5A; 4-cyanobenzenesulfonamide	67	4.49 (6)	566

Example 74

[6-(4-Cyanophenyl)-4-methyl-5-[[[(methylsulfonyl)amino]carbonyl]-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetic acid



45.6 mg (0.08 mmol) of Example 68 are dissolved in 2 ml dichloromethane/trifluoroacetic acid (1:1) and stirred at room temperature for 1 hour. Volatiles are evaporated *in vacuo* and the remainder is purified by RP-HPLC with a water/acetonitrile gradient.

Yield: 13 mg (31% of th.)


- 5 $^1\text{H-NMR}$ (200 MHz, DMSO- d_6): δ = 1.28 (s, 3H), 1.78 (s, 3H), 3.11 (s, 3H), 3.69 (d, 1H), 4.10 (d, 1H), 5.62 (s, 1H), 7.52-8.10 (m, 8H), 11.70 (br s, 1H), 12.70 (br s, 1H) ppm.

In analogy to the procedure for Example 74, the following compounds are prepared:

Example No.	Structure	Starting material	Yield [%]	R _t [min] (method)	Mass [M+H] ⁺
75		Example 69	41	4.56 (1)	605
76		Example 70	73	4.59 (1)	617

Example 77

- 10 *tert.*-Butyl [6-(4-cyanophenyl)-5-(hydroxymethyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate




50 mg (0.13 mmol) of Example 1 are dissolved in 1 ml tetrahydrofuran, 10.8 mg (0.27 mmol) sodium hydride are added, and the mixture is stirred for 15 minutes at room temperature. 27 mg (0.14 mmol) *tert.*-butyl bromoacetate is added, and the mixture is stirred at room temperature for 1
 5 hour. Saturated aq. ammonium chloride solution is added, and the reaction mixture is diluted with ethyl acetate. The organic phase is washed with water, dried over magnesium sulfate and evaporated to dryness. The crude product is purified by column chromatography over silica gel (eluent: cyclohexane/ethyl acetate).

Yield: 17.7 mg (27% of th.)

10 ¹H-NMR (200 MHz, DMSO-d₆): δ = 1.30 (s, 9H), 1.56 (s, 3H), 3.52 (dd, 1H), 3.62 (dd, 1H), 3.92-4.12 (m, 2H), 4.82 (t, 1H), 5.25 (s, 1H), 7.52-7.80 (m, 6H), 7.88 (d, 2H) ppm.

Example 78

3-{4-(4-Cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl}propanoic acid




45.0 mg (0.057 mmol) of Example 17A und 5.00 mg palladium on charcoal (10%) are suspended in 5 ml ethanol and hydrogenated under 1 atm hydrogen at room temperature for 12 minutes. The reaction mixture is filtered over celite and the residue is washed with ethanol. The filtrate is
5 evaporated to dryness *in vacuo* and the remainder is heated without solvent at 130°C under an argon atmosphere for 20 minutes. The crude product is purified by RP-HPLC with a water/acetonitrile gradient.

Yield: 9 mg (26% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 1.53 (s, 3H), 1.92-2.42 (m, 4H), 4.93 (s, 1H), 7.50-7.94 (m,
10 9H), 12.08 (br s, 1H) ppm.

Example 79

((4*R*)-4-(4-Cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl)methylmalonic acid




200 mg (0.25 mmol) of Example 19A und 5.0 mg palladium on charcoal (10%) are suspended in 10 ml ethanol and hydrogenated under 1 atm hydrogen at room temperature for 15 minutes. The reaction mixture is filtered over celite and the residue is washed with ethanol. The filtrate is
5 evaporated to dryness *in vacuo* and the crude product is purified by RP-HPLC with a water/acetonitrile gradient.

Yield: 87 mg (72% of th.)

$^1\text{H-NMR}$ (300 MHz, DMSO- d_6): δ = 1.5 (s, 3H), 2.2 (dd, 2H), 2.6 (dd, 2H), 3.5 (dd, 1H), 5.0 (d, 1H), 7.5 (m, 2H), 7.6-7.7 (m, 4H), 7.8 (d, 1H), 7.9 (m, 2H), 12.8 (br s, 2H) ppm.

10 Example 80

3-((4R)-4-(4-Cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-5-yl)propanoic acid




40 mg (0.08 mmol) of Example 79 are heated without solvent at 130°C under an argon atmosphere for 20 minutes. The crude product is purified by column chromatography (silica, eluent: dichloromethane/methanol 20:1).

Yield: 36 mg (99% of th.)

- 5 ¹H-NMR (300 MHz, DMSO-d₆): δ = 1.5 (s, 3H), 1.9-2.2 (m, 2H), 2.3-2.4 (m, 2H), 4.9 (d, 1H), 7.50-7.7 (m, 6H), 7.8 (d, 1H), 7.9 (m, 2H), 12.1 (br s, 1H) ppm.

Examples 81 and 82

4-{(4*R*)-5-(1-Hydroxyethyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl} benzonitrile



10

- 300 mg (0.75 mmol) of Example 18A are dissolved in 5 ml tetrahydrofuran. At 0°C, 28.5 mg (0.75 mmol) lithium aluminium hydride (as 1 M solution in tetrahydrofuran) are added slowly. The dark-red solution is stirred at 0°C for 30 minutes, then water is added. The pH is adjusted to 3-4 with 1 N hydrochloric acid. The crude product is extracted with ethyl acetate and purified and separated
15 by column chromatography (silica, eluent: dichloromethane/methanol 40:1).

Example 81 (Diastereomer I):

Yield: 15 mg (5% of th.)

¹H-NMR (400 MHz, DMSO-d₆): δ = 1.2 (d, 3H), 1.6 (s, 3H), 4.6 (m, 2H), 4.9 (d, 1H), 7.4 (m, 1H), 7.5 (m, 1H), 7.7 (m, 4H), 7.8 (m, 2H), 7.9 (d, 1H) ppm.


Example 82 (Diastereomer II):

Yield: 15 mg (5% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 0.8 (d, 3H), 1.6 (s, 3H), 4.6 (m, 1H), 5.0 (d, 1H), 5.1 (d, 1H), 7.5 (m, 1H), 7.6-7.7 (m, 5H), 7.8 (d, 1H), 7.9 (m, 2H) ppm.

5 Example 83

4-{5-(4-Bromobenzoyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl}benzonitrile




4.23 g (20.74 mmol) *N*-[3-(trifluoromethyl)phenyl]urea, 2.72 g (20.74 mmol) 4-cyanobenzaldehyde, 5.00 g (20.74 mmol) 1-(4-bromophenyl)butane-1,3-dione and 6.5 g polyphosphoric acid ethyl ester are suspended in 50 ml of tetrahydrofuran. The mixture is stirred at reflux for 20 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate mixtures as eluent.

Yield: 5.32 g (45% of th.)

15 ¹H-NMR (400 MHz, DMSO-d₆): δ = 1.4 (s, 3H), 5.4 (d, 1H), 7.6-7.9 (m, 12H), 8.4 (d, 1H) ppm.

Example 84

tert.-Butyl [5-(4-bromobenzoyl)-6-(4-cyanophenyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate




5.32 g (9.85 mmol) of Example 83 are dissolved in 150 ml tetrahydrofuran, and 0.98 g (24.61 mmol) sodium hydride (60% suspension in mineral oil) are added slowly. After stirring for one hour at room temperature, 2.88 g (14.77 mmol) *tert.*-butyl bromoacetate are added. After stirring at room temperature for one hour, the mixture is quenched with water, the solvent is removed *in vacuo* and the residue is purified by column chromatography (silica, eluent: cyclohexane/ethyl acetate 10:1, 5:1).

Yield: 3.12 g (48% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 1.3 (s, 9H), 1.4 (s, 3H), 3.9 (d, 1H), 4.1 (d, 1H), 5.6 (s, 1H), 7.7 (m, 7H), 7.8 (m, 5H) ppm.

Example 85

[5-(4-Bromobenzoyl)-6-(4-cyanophenyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetic acid




100 mg (0.15 mmol) of Example 84 are dissolved in 10 ml dichloromethane and 0.25 ml trifluoroacetic acid, and the mixture is stirred at room temperature overnight. Volatiles are evaporated *in vacuo* and the remainder is purified by RP-HPLC using a water/acetonitrile gradient.

5 Yield: 47 mg (52% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 1.4 (s, 3H), 3.7 (d, 1H), 4.2 (d, 1H), 5.7 (s, 1H), 7.7 (m, 8H), 7.8 (m, 4H) ppm.

Example 86

10 *tert.*-Butyl [5-(biphenyl-4-ylcarbonyl)-6-(4-cyanophenyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)-phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate




100 mg (0.15 mmol) of Example 84 are dissolved in 5 ml *N,N*-dimethylformamide under an argon atmosphere, and 19 mg (0.15 mmol) phenylboronic acid, 100 mg (0.31 mmol) cesium carbonate and 5 mg dichloro[bis(triphenylphosphino)]palladium are added. The reaction mixture is stirred at 120°C for 18 hours. The product is purified by RP-HPLC using a water/acetonitrile gradient.

5 Yield: 39 mg (39% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 1.3 (s, 9H), 1.5 (s, 3H), 3.9 (d, 1H), 4.2 (d, 1H), 5.7 (s, 1H), 7.5 (m, 3H), 7.7-7.9 (m, 14H) ppm.

Example 87

10 [5-(Biphenyl-4-ylcarbonyl)-6-(4-cyanophenyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetic acid




30 mg (0.05 mmol) of Example 86 are dissolved in 2 ml dichloromethane and 0.07 ml trifluoroacetic acid, and the mixture is stirred at room temperature overnight. Volatiles are evaporated *in vacuo* and the remainder is purified by RP-HPLC using a water/acetonitrile gradient.

15 Yield: 23 mg (83% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 1.5 (s, 3H), 3.7 (d, 1H), 4.3 (d, 1H), 5.7 (s, 1H), 7.4-7.5 (m, 3H), 7.6-7.9 (m, 14H) ppm.

Example 88

20 4-{5-(4-Nitrobenzoyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl}benzonitrile




9.85 g (48.27 mmol) *N*-[3-(trifluoromethyl)phenyl]urea, 6.33 g (48.27 mmol) 4-cyanobenzaldehyde, 10.00 g (48.27 mmol) 1-(4-nitrophenyl)butane-1,3-dione and 15 g polyphosphoric acid ethyl ester are suspended in 100 ml of tetrahydrofuran. The mixture is stirred at reflux for 20
 5 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate mixtures as eluent.

Yield: 7.84 g (32% of th.)

¹H-NMR (400 MHz, DMSO-d₆): δ = 1.4 (s, 3H), 5.4 (d, 1H), 7.6 (m, 3H), 7.7 (m, 1H), 7.8 (m, 2H), 7.8 (m, 1H), 7.9 (m, 3H), 8.3 (m, 2H), 8.5 (d, 1H) ppm.

10 Example 89

tert.-Butyl [5-(4-nitrobenzoyl)-6-(4-cyanophenyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate




5.27 g (10.41 mmol) of Example 88 are dissolved in 150 ml tetrahydrofuran, and 1.04 g (26.02 mmol) sodium hydride (60% suspension in mineral oil) are added slowly. After stirring for one hour at room temperature, 3.04 g (15.61 mmol) *tert.*-butyl bromoacetate are added. After stirring at room temperature for one hour, the mixture is quenched with water, the solvent is removed *in vacuo* and the residue is purified by column chromatography (silica, eluent: cyclohexane/ethyl acetate 10:1, 5:1).

Yield: 0.16 g (3% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 1.3 (s, 9H), 1.5 (s, 3H), 3.9 (d, 1H), 4.2 (d, 1H), 5.6 (s, 1H), 7.7 (m, 4H), 7.8-7.9 (m, 6H), 8.3 (m, 2H) ppm.

10 Example 90

tert.-Butyl [5-(4-aminobenzoyl)-6-(4-cyanophenyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate




160 mg (0.26 mmol) of Example 89 und 5 mg palladium on charcoal (10%) are suspended in 20 ml tetrahydrofuran and hydrogenated under 1 atm hydrogen at room temperature for 18 hours. The reaction mixture is filtered over celite, the filtrate is evaporated to dryness *in vacuo* and the crude product is purified by RP-HPLC with a water/acetonitrile gradient.

Yield: 69 mg (45% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 1.3 (s, 9H), 1.4 (s, 3H), 3.7 (d, 1H), 4.1 (d, 1H), 5.6 (s, 1H), 6.2 (m, 2H), 6.5 (m, 2H), 7.5 (m, 2H), 7.6 (m, 2H), 7.7 (m, 2H), 7.8 (m, 4H) ppm.

Example 91

tert.-Butyl [(6*R*)-6-(4-cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate




- 5 The title compound is prepared from Example 23 according to the procedure described for Example 26.

Yield: 108 mg (85% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 7.90-7.58 (m, 8H), 5.74-5.67 (m, 1H), 4.15-3.85 (m, 2H), 2.30-2.19 (m, 1H), 1.97 (s, 3H), 1.30 (s, 9H), 0.90-0.73 (m, 4H) ppm.

10 *Example 92*

[(6*R*)-6-(4-Cyanophenyl)-5-(cyclopropylcarbonyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetic acid




The title compound is prepared from Example 91 according to the procedure described for Example 30.

Yield: 80 mg (89% of th.)

- 5 $^1\text{H-NMR}$ (300 MHz, DMSO-d_6): δ = 12.62 (br s, 1H), 7.91-7.55 (m, 8H), 5.77 (s, 1H), 4.23-3.70 (m, 2H), 2.32-2.19 (m, 1H), 1.97 (s, 3H), 0.94-0.72 (m, 4H) ppm.

Example 93

4-{5-[(2,2-Dimethyl-1,3-dioxolan-4-yl)carbonyl]-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl}benzonitrile



10

To a stirred solution of Example 20A (120 mg, 0.64 mmol), *N*-[3-(trifluoromethyl)phenyl]urea (109 mg, 0.54 mmol) and 4-cyanobenzaldehyde (84.5 mg, 0.64 mmol) in methyl *tert.*-butyl ether (5


ml) is added 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (683 mg, 1 mmol). The solution is refluxed overnight under argon, then cooled to room temperature, quenched with water (200 ml) and extracted with methyl *tert.*-butyl ether (3 x 100 ml). The combined organic phases are washed with brine, dried over anhydrous magnesium sulfate, filtered and concentrated. The residue is purified by preparative HPLC.

Yield: 15 mg (5% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 8.48 (m, 1H), 8.00-7.38 (m, 8H), 5.66-5.40 (m, 1H), 4.99-4.74 (m, 1H), 4.16-3.85 (m, 2H), 1.94 (d, 3H), 1.35-1.11 (m, 6H) ppm.

Example 94

10 4-{5-(Cyclohexylcarbonyl)-3-[2-(diethylamino)ethyl]-6-methyl-2-oxo-1-[3-(trifluoromethyl)-phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl} benzonitrile




To a stirred solution of Example 21 (780 mg, 1.67 mmol) in THF (25 ml) is added sodium hydride (60% dispersion in mineral oil; 653 mg, 2.53 mmol). The mixture is stirred for 1 hour at room temperature, then 2-bromo-*N,N*-diethylethanamine (653 mg, 2.53 mmol) is added, and the solution is stirred overnight (16 h) at room temperature. The reaction mixture is quenched with methanol (10 ml), concentrated, and the crude product is purified by preparative HPLC.

Yield: 175 mg (18% of th.)

¹H-NMR (400 MHz, DMSO-d₆): δ = 7.98-7.50 (m, 8H), 5.71 (s, 1H), 3.66-3.54 (m, 2H), 3.10-3.00 (m, 2H), 2.86-2.76 (m, 1H), 2.69-2.30 (m, 4H), 1.77 (s, 3H), 1.73-1.00 (m, 10H), 0.93 (t, 6H) ppm.

Example 95

4-{5-(Cyclopropylcarbonyl)-3-[2-(diethylamino)ethyl]-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl} benzonitrile




- 5 The title compound is prepared from Example 22 according to the procedure described for Example 94.

Yield: 235 mg (59% of th.)

¹H-NMR (400 MHz, DMSO-d₆): δ = 7.93-7.54 (m, 8H), 5.81 (s, 1H), 3.68-3.59 (m, 2H), 3.12-3.01 (m, 2H), 2.60-2.56 (m, 5H), 1.92 (s, 3H), 0.97-0.78 (m, 10H) ppm.

10 *Example 96*

tert.-Butyl [6-(4-cyanophenyl)-5-[(2,2-dimethyl-1,3-dioxolan-4-yl)carbonyl]-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2*H*)-yl]acetate




The title compound is prepared from Example 93 according to the procedure described for Example 26.

Yield: 145 mg (62% of th.)

- 5 ¹H-NMR (400 MHz, DMSO-d₆): δ = 7.92-7.56 (m, 8H), 5.77 and 5.62 (s, 1H; diastereomers A and B), 4.97 and 4.82 (t, 1H; diastereomers A and B), 4.19-3.83 (m, 4H), 1.96 and 1.93 (s, 3H; diastereomers A and B), 1.34-1.20 (m, 15H) ppm.

Example 97

- 10 Methyl [6-(4-cyanophenyl)-5-[(2,2-dimethyl-1,3-dioxolan-4-yl)carbonyl]-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydropyrimidin-1(2H)-yl]acetate




The title compound is prepared from Example 93 according to the procedure described for Example 26.

Yield: 77 mg (65% of th.)

¹H-NMR (300 MHz, DMSO-d₆): δ = 7.94-7.55 (m, 8H), 5.81 and 5.66 (s, 1H; 2 diastereomers),
5 4.99 and 4.87 (t, 1H; 2 diastereomers), 4.35-3.85 (m, 4H), 3.57 and 3.55 (s, 3H; 2 diastereomers),
1.95 and 1.92 (s, 3H; 2 diastereomers), 1.25 (dd, 6H) ppm.

Example 98

4-{{(4*R*)-5-(Cyclopropylcarbonyl)-3-[2-(diethylamino)ethyl]-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl} benzonitrile




10

The enantiomers of Example 95 are separated by preparative HPLC on a chiral phase [Daicel Chiralpak AD-H, 5 μm; 250 mm x 20 mm; eluent: 80:20 isohexane/isopropanol with 0.2% diethylamine; flow 15 ml/min; temperature 25°C; detection 220 nm].

R_t = 4.58 min. [Daicel Chiralpak AD-H, 5 μm; 250 mm x 4.6 mm; eluent: 80:20 isohexane/isopropanol with 0.2% diethylamine; flow 1.0 ml/min; temperature 35°C; detection 220 nm].
15

Example 99

4-{{(4*R*)-5-(Cyclohexylcarbonyl)-3-[2-(diethylamino)ethyl]-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidin-4-yl} benzonitrile



The enantiomers of Example 94 are separated by preparative HPLC on a chiral phase [Daicel Chiralpak AD-H, 5 μ m; 250 mm x 20 mm; eluent: 80:20 isohexane/isopropanol with 0.2% diethylamine; flow 15 ml/min; temperature 25°C; detection 220 nm].

- 5 R_t = 4.24 min. [Daicel Chiralpak AD-H, 5 μ m; 250 mm x 4.6 mm; eluent: 80:20 isohexane/isopropanol with 0.2% diethylamine; flow 1.0 ml/min; temperature 35°C; detection 220 nm].

C. Operative examples relating to pharmaceutical compositions

The compounds according to the invention can be converted into pharmaceutical preparations as follows:

Tablet**5 Composition:**

100 mg of the compound of Example 1, 50 mg of lactose (monohydrate), 50 mg of maize starch (native), 10 mg of polyvinylpyrrolidone (PVP 25) (from BASF, Ludwigshafen, Germany) and 2 mg of magnesium stearate.

Tablet weight 212 mg, diameter 8 mm, curvature radius 12 mm.

10 Preparation:

The mixture of active component, lactose and starch is granulated with a 5% solution (m/m) of the PVP in water. After drying, the granules are mixed with magnesium stearate for 5 min. This mixture is moulded using a customary tablet press (tablet format, see above). The moulding force applied is typically 15 kN.

15 Orally administrable suspension**Composition:**

1000 mg of the compound of Example 1, 1000 mg of ethanol (96%), 400 mg of Rhodigel (xanthan gum from FMC, Pennsylvania, USA) and 99 g of water.


20 A single dose of 100 mg of the compound according to the invention is provided by 10 ml of oral suspension.

Preparation:

The Rhodigel is suspended in ethanol and the active component is added to the suspension. The water is added with stirring. Stirring is continued for about 6h until the swelling of the Rhodigel is complete.

We claim

1. Compounds of the general formula (I)



wherein

- 5 A represents an aryl or heteroaryl ring,

R^1 , R^2 and R^3 independently from each other represent hydrogen, halogen, nitro, cyano, C_1 - C_6 -alkyl, hydroxy or C_1 - C_6 -alkoxy, wherein C_1 - C_6 -alkyl and C_1 - C_6 -alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and C_1 - C_4 -alkoxy,

- 10 R^4 represents

- C_1 - C_6 -alkyl which can be substituted by up to three radicals independently selected from the group consisting of hydroxy, C_1 - C_6 -alkoxycarbonyl and hydroxycarbonyl,
- C_3 - C_8 -cycloalkylcarbonyl which can be substituted by up to three radicals independently selected from the group consisting of C_1 - C_6 -alkyl, hydroxy, oxo, C_1 - C_6 -alkoxycarbonyl and hydroxycarbonyl,
- C_1 - C_6 -alkylcarbonyl which is substituted by phenyl- C_1 - C_6 -alkoxy or phenyl- C_1 - C_6 -alkoxycarbonyl which for their part, in the phenyl moiety, can be substituted by halogen, C_1 - C_6 -alkyl, hydroxy, C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxycarbonyl or hydroxycarbonyl,
- C_6 - C_{10} -arylcarbonyl which is substituted by one, two or three radicals independently selected from the group consisting of halogen, cyano, nitro, C_1 - C_6 -

alkyl, trifluoromethyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, amino, C₁-C₆-alkoxycarbonyl, hydroxycarbonyl and phenyl,


- 5 - C₁-C₆-alkoxycarbonyl which is substituted by one or two radicals independently selected from the group consisting of phenyl-C₁-C₆-alkoxy, phenyl-C₁-C₆-alkoxycarbonyl, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonylamino and 5- or 6-membered heterocyclyl, wherein C₁-C₆-alkoxy is further substituted by C₁-C₆-alkoxycarbonyl or hydroxycarbonyl, and 5- or 6-membered heterocyclyl is further substituted by hydroxy, oxo, C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,
- 10 - heteroarylcarbonyl which is substituted by one or two radicals independently selected from the group consisting of hydroxy, amino, halogen, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl, and which can additionally be substituted by C₁-C₆-alkyl,
- 15 - mono- or di-C₁-C₆-alkylaminocarbonyl wherein the alkyl moiety or at least one alkyl moiety, respectively, is substituted by C₆-C₁₀-aryl which for its part can be further substituted by up to three radicals independently selected from the group consisting of halogen, cyano, trifluoromethyl, C₁-C₆-alkyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- 20 - C₆-C₁₀-arylamino carbonyl or *N*-(C₁-C₆-alkyl)-*N*-(C₆-C₁₀-aryl)aminocarbonyl wherein aryl is substituted by one, two or three radicals independently selected from the group consisting of halogen, cyano, trifluoromethyl, C₁-C₆-alkyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl, and wherein alkyl, when present, can be substituted by up to three radicals independently selected from the group consisting of hydroxy, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- 25 - C₃-C₈-cycloalkylaminocarbonyl or *N*-(C₁-C₆-alkyl)-*N*-(C₃-C₈-cycloalkyl)aminocarbonyl wherein cycloalkyl can be substituted by up to three radicals independently selected from the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl, and wherein alkyl, when present, can be substituted by up to three radicals independently selected from the group consisting of hydroxy, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- 30 - heterocyclylcarbonyl which is substituted by one, two or three radicals independently selected from the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-

- 5 C₆-alkoxy, C₁-C₆-alkoxycarbonyl, phenyl-C₁-C₆-alkoxycarbonyl, hydroxycarbonyl, 5- or 6-membered heterocyclyl, 5- or 6-membered heteroaryl and C₆-C₁₀-aryl, wherein C₁-C₆-alkyl is further substituted by hydroxy, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl or hydroxycarbonyl, and wherein C₆-C₁₀-aryl can be further substituted by up to three radicals independently selected from the group consisting of halogen, cyano, trifluoromethyl, C₁-C₆-alkyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- 10 – N-(heterocyclyl)aminocarbonyl wherein heterocyclyl can be further substituted by up to three radicals independently selected from the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl, hydroxycarbonyl and phenyl-C₁-C₆-alkyl,
- 15 – a group of the formula -C(=O)-NR^a-SO₂-R^b wherein R^a represents hydrogen or C₁-C₆-alkyl, and R^b represents C₁-C₆-alkyl which can be substituted by trifluoromethyl, or R^b represents C₆-C₁₀-aryl which can be substituted by C₁-C₆-alkyl, halogen, cyano, nitro or trifluoromethyl,
- or
- a group of the formula -P(=O)(OR^c)₂ wherein R^c represents hydrogen or C₁-C₆-alkyl,
- 20 R⁵ represents C₁-C₄-alkyl, which can be substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy, C₁-C₆-alkoxy, C₂-C₆-alkenoxy, C₁-C₆-alkylthio, amino, mono- and di-C₁-C₆-alkylamino, arylamino, hydroxycarbonyl, C₁-C₆-alkoxycarbonyl and the radical -O-C₁-C₄-alkyl-O-C₁-C₄-alkyl,
- 25 R⁶ represents hydrogen, C₁-C₆-alkyl, formyl, aminocarbonyl, mono- or di-C₁-C₄-alkylaminocarbonyl, C₃-C₈-cycloalkylcarbonyl, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, N-(C₁-C₄-alkylsulfonyl)-aminocarbonyl, N-(C₁-C₄-alkylsulfonyl)-N-(C₁-C₄-alkyl)-aminocarbonyl, heteroaryl, heterocyclyl, heteroarylcarbonyl or heterocyclylcarbonyl, wherein C₁-C₆-alkyl, mono- and di-C₁-C₄-alkylaminocarbonyl, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, heteroaryl and heterocyclyl can be substituted with one to three identical or different radicals selected from the group consisting of aryl, heteroaryl, hydroxy, C₁-C₄-alkoxy, hydroxycarbonyl, C₁-C₆-alkoxycarbonyl, aminocarbonyl, mono- and di-C₁-C₄-alkylaminocarbonyl, amino,
- 30

mono- and di-C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, tri-(C₁-C₆-alkyl)-silyl, cyano, *N*-(mono- or di-C₁-C₄-alkylamino-C₁-C₄-alkyl)-aminocarbonyl, *N*-(C₁-C₄-alkoxy-C₁-C₄-alkyl)-aminocarbonyl and halogen,

or

5 R⁶ represents a moiety of the formula



wherein

R^d is selected from the group consisting of hydrogen and C₁-C₆-alkyl, and

n represents an integer of 1 or 2,

10 or

R⁶ represents a group of the formula -T-U wherein

T represents a C₁-C₆-alkanediyl or C₂-C₆-alkenediyl group

and

U represents

- 15
- C₆-C₁₀-aryl or 5- or 6-membered heteroaryl each of which is substituted by one, two or three radicals independently selected from the group consisting of halogen, C₁-C₆-alkyl, 5- or 6-membered heteroaryl and a group of the formula -V-W wherein V represents a bond or a C₁-C₆-alkanediyl or C₂-C₆-alkenediyl group both of which can be further substituted by C₃-C₈-cycloalkyl, and W represents C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,

20

 - a group of the formula -C(=O)-NR^e-SO₂-R^f wherein R^e represents hydrogen or C₁-C₆-alkyl, and R^f represents C₁-C₆-alkyl which can be substituted by trifluoromethyl, or R^f represents C₆-C₁₀-aryl which can be substituted by C₁-C₆-alkyl, halogen, cyano, nitro or trifluoromethyl,

- a group of the formula $-C(=O)-NR^gR^h$ wherein R^g represents hydrogen or C_1-C_6 -alkyl, and R^h represents C_6-C_{10} -aryl which can be substituted by C_1-C_6 -alkoxycarbonyl or hydroxycarbonyl,

5

- a group of the formula $-C(=O)-NR^i-OR^k$ wherein R^i and R^k independently from each other represent hydrogen or C_1-C_6 -alkyl,

or

- C_6-C_{10} -arylalkoxy which, in the aryl part, can be substituted by halogen, C_1-C_6 -alkyl, C_1-C_6 -alkoxycarbonyl or hydroxycarbonyl,

or

10

 R^6 represents

- C_3-C_8 -cycloalkyl which can be substituted by up to three radicals independently selected from the group consisting of C_1-C_6 -alkyl, hydroxy, oxo, C_1-C_6 -alkoxycarbonyl and hydroxycarbonyl,
- C_2-C_6 -alkenyl which can be substituted by C_1-C_6 -alkoxycarbonyl or hydroxycarbonyl,
- C_1-C_6 -alkylcarbonyl which is substituted by C_1-C_6 -alkoxycarbonylamino,
- C_1-C_6 -alkoxycarbonyl which is substituted by phenyl- C_1-C_6 -alkoxycarbonyl which for its part, in the phenyl moiety, can be further substituted by halogen, C_1-C_6 -alkyl, C_1-C_6 -alkoxycarbonyl or hydroxycarbonyl,

15

20

or

- a group of the formula $-SO_2-R^m$ wherein R^m represents C_1-C_6 -alkyl which can be substituted by trifluoromethyl, or R^m represents C_6-C_{10} -aryl which can be substituted by C_1-C_6 -alkyl, halogen, cyano, nitro, trifluoromethyl, C_1-C_6 -alkoxycarbonyl or hydroxycarbonyl,

25

R^7 represents halogen, nitro, cyano, C_1-C_6 -alkyl, hydroxy or C_1-C_6 -alkoxy, wherein C_1-C_6 -alkyl and C_1-C_6 -alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and C_1-C_4 -alkoxy,

and

Y^1 , Y^2 , Y^3 , Y^4 and Y^5 independently from each other represent CH or N, wherein the ring contains either 0, 1 or 2 nitrogen atoms,

and their salts, hydrates and/or solvates and their tautomeric forms.

5 2. Compounds of general formula (I) according to Claim 1, wherein

A represents an aryl or heteroaryl ring,

R^1 , R^2 and R^3 independently from each other represent hydrogen, halogen, nitro, cyano, C_1 - C_6 -alkyl, hydroxy or C_1 - C_6 -alkoxy, wherein C_1 - C_6 -alkyl and C_1 - C_6 -alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and C_1 - C_4 -alkoxy,

10

R^4 represents

- C_1 - C_6 -alkyl which can be substituted by up to three radicals independently selected from the group consisting of hydroxy, C_1 - C_6 -alkoxycarbonyl and hydroxycarbonyl,

- C_3 - C_8 -cycloalkylcarbonyl which can be substituted by up to three radicals independently selected from the group consisting of C_1 - C_6 -alkyl, hydroxy, oxo, C_1 - C_6 -alkoxycarbonyl and hydroxycarbonyl,

15

- C_6 - C_{10} -arylcarbonyl which is substituted by one, two or three radicals independently selected from the group consisting of halogen, cyano, C_1 - C_6 -alkyl, trifluoromethyl, hydroxy, C_1 - C_6 -alkoxy, trifluoromethoxy, C_1 - C_6 -alkoxycarbonyl and hydroxycarbonyl,

20

- C_1 - C_6 -alkoxycarbonyl which is substituted by one or two radicals independently selected from the group consisting of phenyl- C_1 - C_6 -alkoxy, phenyl- C_1 - C_6 -alkoxycarbonyl, C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxycarbonylamino and 5- or 6-membered heterocyclyl, wherein C_1 - C_6 -alkoxy is further substituted by C_1 - C_6 -alkoxycarbonyl or hydroxycarbonyl, and 5- or 6-membered heterocyclyl is further substituted by hydroxy, oxo, C_1 - C_6 -alkoxycarbonyl or hydroxycarbonyl,

25

- heteroarylcarbonyl which is substituted by one or two radicals independently selected from the group consisting of hydroxy, amino, halogen, C_1 - C_6 -alkoxy, C_1 -

C₆-alkoxycarbonyl and hydroxycarbonyl, and which can additionally be substituted by C₁-C₆-alkyl,

- 5 - mono- or di-C₁-C₆-alkylaminocarbonyl wherein the alkyl moiety or at least one alkyl moiety, respectively, is substituted by C₆-C₁₀-aryl which for its part can be further substituted by up to three radicals independently selected from the group consisting of halogen, cyano, trifluoromethyl, C₁-C₆-alkyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,
- 10 - heterocyclylcarbonyl which is substituted by one, two or three radicals independently selected from the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl, phenyl-C₁-C₆-alkoxycarbonyl, hydroxycarbonyl, 5- or 6-membered heterocyclyl, 5- or 6-membered heteroaryl and C₆-C₁₀-aryl, wherein C₁-C₆-alkyl is further substituted by hydroxy, C₁-C₆-alkoxy, C₁-C₆-alkoxycarbonyl or hydroxycarbonyl, and wherein C₆-C₁₀-aryl can be further substituted by up to three radicals independently selected from the group consisting of
- 15 halogen, cyano, trifluoromethyl, C₁-C₆-alkyl, hydroxy, C₁-C₆-alkoxy, trifluoromethoxy, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,

or

- 20 - a group of the formula -C(=O)-NH-SO₂-R^b wherein R^b represents C₁-C₆-alkyl which can be substituted by trifluoromethyl, or R^b represents C₆-C₁₀-aryl which can be substituted by C₁-C₆-alkyl, halogen, cyano, nitro or trifluoromethyl,


25 R⁵ represents C₁-C₄-alkyl, which can be substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy, C₁-C₆-alkoxy, C₂-C₆-alkenoxy, C₁-C₆-alkylthio, amino, mono- and di-C₁-C₆-alkylamino, arylamino, hydroxycarbonyl, C₁-C₆-alkoxycarbonyl and the radical -O-C₁-C₄-alkyl-O-C₁-C₄-alkyl,

30 R⁶ represents hydrogen, C₁-C₆-alkyl, aminocarbonyl, mono- or di-C₁-C₄-alkylaminocarbonyl, C₃-C₈-cycloalkylcarbonyl, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, *N*-(C₁-C₄-alkylsulfonyl)-aminocarbonyl, *N*-(C₁-C₄-alkylsulfonyl)-*N*-(C₁-C₄-alkyl)-aminocarbonyl, heteroarylcarbonyl or heterocyclylcarbonyl, wherein C₁-C₆-alkyl, mono- and di-C₁-C₄-alkylaminocarbonyl, C₁-C₆-alkylcarbonyl and C₁-C₆-alkoxycarbonyl can be substituted with one to three identical or different radicals selected from the group consisting of aryl, heteroaryl, hydroxy, C₁-C₄-alkoxy, hydroxy-

carbonyl, C₁-C₆-alkoxycarbonyl, aminocarbonyl, mono- and di-C₁-C₄-alkylamino-carbonyl, amino, mono- and di-C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, *N*-(mono- or di-C₁-C₄-alkylamino-C₁-C₄-alkyl)-aminocarbonyl, *N*-(C₁-C₄-alkoxy-C₁-C₄-alkyl)-aminocarbonyl and halogen,

5 or

R⁶ represents a moiety of the formula



wherein

R^d is selected from the group consisting of hydrogen and C₁-C₆-alkyl, and

10 n represents an integer of 1 or 2,

or

R⁶ represents a group of the formula -T-U wherein

T represents a C₁-C₄-alkanediyl or C₂-C₄-alkenediyl group

and

15 U represents

- C₆-C₁₀-aryl or 5- or 6-membered heteroaryl each of which is substituted by one, two or three radicals independently selected from the group consisting of halogen, C₁-C₆-alkyl, 5- or 6-membered heteroaryl and a group of the formula -V-W wherein V represents a bond, a C₂-C₆-alkenediyl group or a C₁-C₆-alkanediyl group the latter of which can be further substituted by C₃-C₈-cycloalkyl, and W represents C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,
- a group of the formula -C(=O)-NH-SO₂-R^f wherein R^f represents C₁-C₆-alkyl which can be substituted by trifluoromethyl, or R^f represents C₆-C₁₀-

20

aryl which can be substituted by C₁-C₆-alkyl, halogen, cyano, nitro or trifluoromethyl,

or

- a group of the formula -C(=O)-NHR^h wherein R^h represents C₆-C₁₀-aryl which can be substituted by C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,

or

R⁶ represents

- C₃-C₈-cycloalkyl which can be substituted by up to three radicals independently selected from the group consisting of C₁-C₆-alkyl, hydroxy, oxo, C₁-C₆-alkoxycarbonyl and hydroxycarbonyl,

or

- C₂-C₆-alkenyl which can be substituted by C₁-C₆-alkoxycarbonyl or hydroxycarbonyl,

R⁷ represents halogen, nitro, cyano, C₁-C₆-alkyl, hydroxy or C₁-C₆-alkoxy, wherein C₁-C₆-alkyl and C₁-C₆-alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and C₁-C₄-alkoxy,

and

Y¹, Y², Y³, Y⁴ and Y⁵ independently from each other represent CH or N, wherein the ring contains either 0, 1 or 2 nitrogen atoms.

3. Compounds of general formula (I) according to Claim 1, wherein

A represents a phenyl, naphthyl or pyridyl ring,

R¹, R² and R³ independently from each other represent hydrogen, fluoro, chloro, bromo, nitro, cyano, methyl, ethyl, trifluoromethyl or trifluoromethoxy,

R⁴ represents

- C₁-C₄-alkyl which can be substituted by up to two radicals independently selected from the group consisting of hydroxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- C₃-C₆-cycloalkylcarbonyl which can be substituted by up to two radicals independently selected from the group consisting of C₁-C₄-alkyl, hydroxy, oxo, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- benzoyl which is substituted by one, two or three radicals independently selected from the group consisting of fluoro, chloro, bromo, cyano, C₁-C₄-alkyl, trifluoromethyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- C₁-C₄-alkoxycarbonyl which is substituted by one or two radicals independently selected from the group consisting of benzyloxy, benzyloxycarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonylamino, pyrrolidinyl, piperidinyl and morpholinyl, wherein C₁-C₄-alkoxy is further substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl, and wherein pyrrolidinyl, piperidinyl and morpholinyl is further substituted by hydroxy, oxo, C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,
- furylcarbonyl, thienylcarbonyl, oxazolylcarbonyl, thiazolylcarbonyl, pyridylcarbonyl or pyrimidinylcarbonyl each of which is substituted by one or two radicals independently selected from the group consisting of hydroxy, amino, fluoro, chloro, bromo, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl, and each of which can additionally be substituted by C₁-C₄-alkyl,
- mono- or di-C₁-C₄-alkylaminocarbonyl wherein the alkyl moiety or at least one alkyl moiety, respectively, is substituted by phenyl which for its part can be further substituted by up to three radicals independently selected from the group consisting of fluoro, chloro, bromo, cyano, trifluoromethyl, C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- tetrahydrofurylcarbonyl, tetrahydropyranylcabonyl, piperidinylcarbonyl, piperazinylcarbonyl or morpholinylcarbonyl each of which is substituted by one or two radicals independently selected from the group consisting of C₁-C₄-alkyl, hydroxy, oxo, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl, benzyloxycarbonyl, hydroxycarbonyl, piperidinyl, morpholinyl, pyridyl and phenyl, wherein C₁-C₄-alkyl is further substituted by hydroxy, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl or hydroxycarbonyl, and wherein phenyl can be further substituted by up to three radicals independently

selected from the group consisting of fluoro, chloro, bromo, cyano, trifluoromethyl, C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,

or


- 5 – a group of the formula $-C(=O)-NH-SO_2-R^b$ wherein R^b represents C₁-C₄-alkyl which can be substituted by trifluoromethyl, or R^b represents phenyl which can be substituted by C₁-C₄-alkyl, fluoro, chloro, bromo, cyano, nitro or trifluoromethyl,

R⁵ represents methyl or ethyl,

- 10 R⁶ represents hydrogen, C₁-C₆-alkyl, mono- or di-C₁-C₄-alkylaminocarbonyl, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl or heterocyclylcarbonyl, wherein C₁-C₆-alkyl and C₁-C₆-alkoxycarbonyl can be substituted with one to three identical or different radicals selected from the group consisting of hydroxy, C₁-C₄-alkoxy, hydroxycarbonyl, C₁-C₆-alkoxycarbonyl, aminocarbonyl, mono- and di-C₁-C₄-alkylaminocarbonyl, amino, mono- and di-C₁-C₄-alkylamino,

15 or

R⁶ represents a moiety of the formula



wherein

R^d is selected from the group consisting of hydrogen and C₁-C₄-alkyl, and

20 n represents an integer of 1 or 2,

or

R⁶ represents a group of the formula $-T-U$ wherein

T represents a C₁-C₄-alkanediyl group

and

U represents

- phenyl, furyl, thienyl, oxazolyl, thiazolyl or pyridyl each of which is substituted by one or two radicals independently selected from the group consisting of fluoro, chloro, bromo, C₁-C₄-alkyl, thienyl, pyridyl and a group of the formula -V-W wherein V represents a bond or a C₁-C₄-alkanediyl or C₂-C₄-alkenediyl group, and W represents C₁-C₄-alkoxy-carbonyl or hydroxycarbonyl,

- a group of the formula -C(=O)-NH-SO₂-R^f wherein R^f represents C₁-C₄-alkyl which can be substituted by trifluoromethyl, or R^f represents phenyl which can be substituted by C₁-C₄-alkyl, fluoro, chloro, bromo, cyano, nitro or trifluoromethyl,

or

- a group of the formula -C(=O)-NHR^h wherein R^h represents phenyl which can be substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,

or

R⁶ represents

- C₃-C₆-cycloalkyl which can be substituted by up to two radicals independently selected from the group consisting of C₁-C₄-alkyl, hydroxy, oxo, C₁-C₄-alkoxy-carbonyl and hydroxycarbonyl,

or

- C₂-C₄-alkenyl which is substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,

R⁷ represents halogen, nitro, cyano, trifluoromethyl, trifluoromethoxy, methyl or ethyl,

and

Y¹, Y², Y³, Y⁴ and Y⁵ each represent CH.

4. Compounds of general formula (I) according to Claim 1, wherein

A represents a phenyl or a pyridyl ring,

R¹ and R³ each represent hydrogen,

R² represents fluoro, chloro, bromo, nitro or cyano,

R⁴ represents

- 5 – C₁-C₄-alkyl which can be substituted by up to two radicals independently selected from the group consisting of hydroxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- C₃-C₆-cycloalkylcarbonyl which can be substituted by up to two radicals independently selected from the group consisting of C₁-C₄-alkyl, hydroxy, oxo, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- 10 – benzoyl which is substituted by one, two or three radicals independently selected from the group consisting of fluoro, chloro, bromo, cyano, C₁-C₄-alkyl, trifluoromethyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- 15 – C₁-C₄-alkoxycarbonyl which is substituted by one or two radicals independently selected from the group consisting of benzyloxy, benzyloxycarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonylamino, pyrrolidinyl, piperidinyl and morpholinyl, wherein C₁-C₄-alkoxy is further substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl, and wherein pyrrolidinyl, piperidinyl and morpholinyl is further substituted by hydroxy, oxo, C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,
- 20 – furylcarbonyl, oxazolylcarbonyl, thiazolylcarbonyl or pyridylcarbonyl each of which is substituted by one or two radicals independently selected from the group consisting of hydroxy, amino, fluoro, chloro, bromo, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl, and each of which can additionally be substituted by C₁-C₄-alkyl,
- 25 – mono- or di-C₁-C₄-alkylaminocarbonyl wherein the alkyl moiety or at least one alkyl moiety, respectively, is substituted by phenyl which for its part can be further substituted by up to three radicals independently selected from the group consisting of fluoro, chloro, bromo, cyano, trifluoromethyl, C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,
- 30 – piperidinylcarbonyl, piperazinylcarbonyl or morpholinylcarbonyl each of which is substituted by one or two radicals independently selected from the group con-

sisting of C₁-C₄-alkyl, hydroxy, oxo, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl, benzyl-oxycarbonyl, hydroxycarbonyl, piperidiny, morpholinyl, pyridyl and phenyl, wherein C₁-C₄-alkyl is further substituted by hydroxy, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl or hydroxycarbonyl, and wherein phenyl can be further substituted by up to three radicals independently selected from the group consisting of fluoro, chloro, bromo, cyano, trifluoromethyl, C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkoxycarbonyl and hydroxycarbonyl,

or


- a group of the formula -C(=O)-NH-SO₂-R^b wherein R^b represents C₁-C₄-alkyl which can be substituted by trifluoromethyl, or R^b represents phenyl which can be substituted by C₁-C₄-alkyl, fluoro, chloro, bromo, cyano, nitro or trifluoromethyl,

R⁵ represents methyl,

R⁶ represents hydrogen, C₁-C₄-alkyl, mono- or di-C₁-C₄-alkylaminocarbonyl, C₁-C₄-alkylcarbonyl or C₁-C₄-alkoxycarbonyl, wherein C₁-C₄-alkyl and C₁-C₄-alkoxycarbonyl can be substituted with a radical selected from the group consisting of hydroxy, C₁-C₄-alkoxy, C₁-C₄-alkoxycarbonyl, hydroxycarbonyl, aminocarbonyl, mono- and di-C₁-C₄-alkylaminocarbonyl, amino, mono- and di-C₁-C₄-alkylamino,

or

R⁶ represents a moiety of the formula



wherein

R^d is selected from the group consisting of hydrogen and methyl,

or

R⁶ represents a group of the formula -T-U wherein

T represents a -CH₂- group

and

U represents

- phenyl, furyl or oxazolyl each of which is substituted by one or two radicals independently selected from the group consisting of fluoro, chloro, bromo, C₁-C₄-alkyl and a group of the formula -V-W wherein V represents a bond, a -CH₂- group or a -CH=CH- group, and W represents C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,

- a group of the formula -C(=O)-NH-SO₂-R^f wherein R^f represents C₁-C₄-alkyl which can be substituted by trifluoromethyl, or R^f represents phenyl which can be substituted by C₁-C₄-alkyl, fluoro, chloro, bromo, cyano, nitro or trifluoromethyl,

or

- a group of the formula -C(=O)-NHR^h wherein R^h represents phenyl which can be substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,

or

R⁶ represents

- C₃-C₆-cycloalkyl which can be substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,

or

- a -CH=CH- group which is substituted by C₁-C₄-alkoxycarbonyl or hydroxycarbonyl,


R⁷ represents trifluoromethyl or nitro,

and

Y¹, Y², Y³, Y⁴ and Y⁵ each represent CH.

5. Compounds of general formula (I) according to any of the preceding claims, wherein A is phenyl or pyridyl.

6. Compounds of general formula (I) according to any of the preceding claims, wherein R¹ is hydrogen.
7. Compounds of general formula (I) according to any of the preceding claims, wherein R² is cyano.
- 5 8. Compounds of general formula (I) according to any of the preceding claims, wherein R³ is hydrogen.
9. Compounds of general formula (I) according to any of the preceding claims, wherein R⁵ is methyl.
10. Compounds of general formula (I) according to any of the preceding claims, wherein R⁷ is trifluoromethyl or nitro.
- 10 11. Compounds of general formula (IA)




wherein

Z represents CH or N, and

15 R¹, R³, R⁴ and R⁶ have the meaning indicated in any of the preceding claims.


12. Process for synthesizing the compounds of general formula (I) according to Claim 1 by condensing compounds of general formula (II)

136




wherein A, R¹ and R² have the meaning indicated in Claim 1,

with compounds of general formula (III)




5 wherein R⁴ and R⁵ have the meaning indicated in Claim 1,

and compounds of general formula (IV)




wherein R³, R⁷, and Y¹ to Y⁵ have the meaning indicated in Claim 1,

in the presence of an acid or acid anhydride to give compounds of the general formula (IB)



wherein A, R¹ to R⁵, R⁷, and Y¹ to Y⁵ have the meaning indicated in Claim 1,

optionally followed, in case R⁶ does not represent hydrogen, by reaction of the compounds of general formula (IB) with compounds of the general formula (V)



5 wherein

R^{6*} has the meaning of R⁶ as indicated in Claim 1, but does not represent hydrogen, and

X represents a leaving group,

in the presence of a base.

- 10 13. The composition containing at least one compound of general formula (I) according to Claim 1 and a pharmacologically acceptable diluent.
14. A composition according to Claim 13 for the treatment of acute and chronic inflammatory, ischaemic and/or remodelling processes.
- 15 15. The process for the preparation of compositions according to Claim 13 or 14 characterized in that the compounds of general formula (I) according to Claim 1 together with customary auxiliaries are brought into a suitable application form.
16. Use of the compounds of general formula (I) according to Claim 1 for the preparation of medicaments.
- 20 17. Use according to Claim 16 for the preparation of medicaments for the treatment of acute and chronic inflammatory, ischaemic and/or remodelling processes.
18. Use according to Claim 17, wherein the process is chronic obstructive pulmonary disease, acute coronary syndrome, acute myocardial infarction or development of heart failure.
- 25 19. Process for controlling chronic obstructive pulmonary disease, acute coronary syndrome, acute myocardial infarction or development of heart failure in humans and animals by administration of a neutrophil elastase inhibitory amount of at least one compound of general formula (I) according to Claim 1.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2005/001486

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/22 C07D401/06 C07D401/12 C07D403/06 C07D403/12
C07D405/06 C07D413/06 C07D417/06 C07F9/6512 A61K31/505
A61K31/506 A61K31/662 A61P11/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07F A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 03/053930 A (BAYER AKTIENGESellschaft) 3 July 2003 (2003-07-03) the whole document	1, 19
A	OHMOTO K ET AL: "Development of orally active nonpeptidic inhibitors of human neutrophil elastase" JOURNAL OF MEDICINAL CHEMISTRY, vol. 44, no. 8, 12 April 2001 (2001-04-12), pages 1268-1285, XP002908876 the whole document	1, 19
P, X	WO 2004/024700 A (BAYER HEALTHCARE AG) 25 March 2004 (2004-03-25) the whole document	1-19

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

18 July 2005

Date of mailing of the international search report

27/07/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Allard, M

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2005/001486

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 19 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2005/001486

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03053930	A	03-07-2003	GB 2383326 A	25-06-2003
			GB 2390849 A	21-01-2004
			AU 2002361992 A1	09-07-2003
			CA 2470813 A1	03-07-2003
			WO 03053930 A1	03-07-2003
			EP 1458682 A1	22-09-2004
<hr/>				
WO 2004024700	A	25-03-2004	AU 2003282006 A1	30-04-2004
			CA 2498051 A1	25-03-2004
			WO 2004024700 A1	25-03-2004
			EP 1546113 A1	29-06-2005
			GB 2392910 A	17-03-2004
<hr/>				